

BA: 1.3

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.
- (vi) $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.
- (vi) $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$.

Proof: (i):

Case $x \geq 0$:

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.
- (vi) $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$.

Proof: (i):

Case $x \geq 0$: $|x| = x \geq 0$

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.
- (vi) $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$.

Proof: (i):

Case $x \geq 0$: $|x| = x \geq 0$ also $|x| = x = 0 \Leftrightarrow x = 0$.

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.
- (vi) $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$.

Proof: (i):

Case $x \geq 0$: $|x| = x \geq 0$ also $|x| = x = 0 \Leftrightarrow x = 0$.

Case $x < 0$: $|x| = -x > 0$

The *absolute value* is the size of $x \in \mathbb{R}$:

$$|x| := \begin{cases} x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

Proposition

- (i) $|x| \geq 0$, moreover, $|x| = 0 \Leftrightarrow x = 0$.
- (ii) $|-x| = |x|$ for all $x \in \mathbb{R}$.
- (iii) $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$.
- (iv) $|x|^2 = x^2$ for all $x \in \mathbb{R}$.
- (v) $|x| \leq y \Leftrightarrow -y \leq x \leq y$.
- (vi) $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$.

Proof: (i):

Case $x \geq 0$: $|x| = x \geq 0$ also $|x| = x = 0 \Leftrightarrow x = 0$.

Case $x < 0$: $|x| = -x > 0$ and $|x| \neq 0$.

(ii) “ $|-x| = |x|$ for all $x \in \mathbb{R}$ ”:

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$:

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0$

(ii) “ $|-x| = |x|$ for all $x \in \mathbb{R}$ ”:

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

(ii) “ $|-x| = |x|$ for all $x \in \mathbb{R}$ ”:

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(ii) “ $|-x| = |x|$ for all $x \in \mathbb{R}$ ”:

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) “ $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ”:

(ii) “ $|-x| = |x|$ for all $x \in \mathbb{R}$ ”:

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) “ $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ”:

Case $x = 0$ or $y = 0$: immediate.

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$:

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0$

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy$

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$:

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0$

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$:

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0$

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0 \Rightarrow |xy| = -(xy) = x(-y) = |x| |y|$.

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0 \Rightarrow |xy| = -(xy) = x(-y) = |x| |y|$.

Case $x < 0$ & $y > 0$ is similar.

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0 \Rightarrow |xy| = -(xy) = x(-y) = |x| |y|$.

Case $x < 0$ & $y > 0$ is similar.

(iv) " $|x|^2 = x^2$ for all $x \in \mathbb{R}$ ":

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0 \Rightarrow |xy| = -(xy) = x(-y) = |x| |y|$.

Case $x < 0$ & $y > 0$ is similar.

(iv) " $|x|^2 = x^2$ for all $x \in \mathbb{R}$ ":

Case $x \geq 0$: immediate.

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0 \Rightarrow |xy| = -(xy) = x(-y) = |x| |y|$.

Case $x < 0$ & $y > 0$ is similar.

(iv) " $|x|^2 = x^2$ for all $x \in \mathbb{R}$ ":

Case $x \geq 0$: immediate.

Case $x < 0$:

(ii) " $|-x| = |x|$ for all $x \in \mathbb{R}$ ":

Case $x > 0$: $-x < 0 \Rightarrow |-x| = -(-x) = x = |x|$.

Cases $x < 0$ and $x = 0$ similar.

(iii) " $|xy| = |x| |y|$ for all $x, y \in \mathbb{R}$ ":

Case $x = 0$ or $y = 0$: immediate.

Case $x > 0$ & $y > 0$: $xy > 0 \Rightarrow |xy| = xy = |x| |y|$.

Case $x < 0$ & $y < 0$: $xy = (-x)(-y) > 0 \Rightarrow |xy| = xy = (-x)(-y) = |x| |y|$.

Case $x > 0$ & $y < 0$: $xy < 0 \Rightarrow |xy| = -(xy) = x(-y) = |x| |y|$.

Case $x < 0$ & $y > 0$ is similar.

(iv) " $|x|^2 = x^2$ for all $x \in \mathbb{R}$ ":

Case $x \geq 0$: immediate.

Case $x < 0$: $|x|^2 = (-x)^2 = x^2$.

$$(v) \text{ “} |x| \leq y \iff -y \leq x \leq y \text{”}:$$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$:

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$:

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$:

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

Case $x < 0$:

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

Case $x < 0$: $-y \leq x$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

Case $x < 0$: $-y \leq x \Rightarrow y \geq -x$

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

Case $x < 0$: $-y \leq x \Rightarrow y \geq -x = |x|$.

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

Case $x < 0$: $-y \leq x \Rightarrow y \geq -x = |x|$.

(vi) " $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$ ":

(v) " $|x| \leq y \Leftrightarrow -y \leq x \leq y$ ":

Suppose $|x| \leq y$. As $|x| \geq 0 \Rightarrow y \geq 0$.

Case $x \geq 0$: $x \leq y$, $y \geq 0 \Rightarrow -y \leq 0 \leq x \Rightarrow -y \leq x \leq y$.

Case $x < 0$: $-x = |x| \leq y \Rightarrow x \geq -y$, $y \geq 0 > x \Rightarrow -y \leq x \leq y$.

Now suppose $-y \leq x \leq y$.

Case $x \geq 0$: $y \geq x = |x|$.

Case $x < 0$: $-y \leq x \Rightarrow y \geq -x = |x|$.

(vi) " $-|x| \leq x \leq |x|$ for all $x \in \mathbb{R}$ ":

(v) with $y = |x|$.



Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof: By the previous proposition $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$.

Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof: By the previous proposition $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$.

Add the two: $-(|x| + |y|) \leq x + y \leq |x| + |y|$.

Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof: By the previous proposition $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$.

Add the two: $-(|x| + |y|) \leq x + y \leq |x| + |y|$.

By the proposition again $|x + y| \leq |x| + |y|$.



Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof: By the previous proposition $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$.

Add the two: $-(|x| + |y|) \leq x + y \leq |x| + |y|$.

By the proposition again $|x + y| \leq |x| + |y|$.



E.g.,

Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof: By the previous proposition $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$.

Add the two: $-(|x| + |y|) \leq x + y \leq |x| + |y|$.

By the proposition again $|x + y| \leq |x| + |y|$. □

E.g.,

$$3 = |1 + 2| \leq |1| + |2| = 1 + 2$$

Proposition (Triangle Inequality)

$|x + y| \leq |x| + |y|$ for all $x, y \in \mathbb{R}$.

Proof: By the previous proposition $-|x| \leq x \leq |x|$ and $-|y| \leq y \leq |y|$.

Add the two: $-(|x| + |y|) \leq x + y \leq |x| + |y|$.

By the proposition again $|x + y| \leq |x| + |y|$. □

E.g.,

$$3 = |1 + 2| \leq |1| + |2| = 1 + 2$$

$$1 = |1 - 2| = |1 + (-2)| \leq |1| + |-2| = 1 + 2 = 3$$

Corollary

Let $x, y \in \mathbb{R}$.

(i) (*reverse triangle inequality*) $\big||x| - |y|\big| \leq |x - y|.$

Corollary

Let $x, y \in \mathbb{R}$.

- (i) (*reverse triangle inequality*) $\big||x| - |y|\big| \leq |x - y|$.
- (ii) $|x - y| \leq |x| + |y|$.

Corollary

Let $x, y \in \mathbb{R}$.

(i) (*reverse triangle inequality*) $\big||x| - |y|\big| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b|$$

Corollary

Let $x, y \in \mathbb{R}$.

(i) (*reverse triangle inequality*) $\big||x| - |y|\big| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b| \quad \Rightarrow \quad |a| - |b| \leq |a - b|$$

Corollary

Let $x, y \in \mathbb{R}$.

(i) (reverse triangle inequality) $\left|(|x| - |y|)\right| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b| \quad \Rightarrow \quad |a| - |b| \leq |a - b|$$

Switch roles of a and b :

$$|b| - |a| \leq |b - a| = |a - b|$$

Corollary

Let $x, y \in \mathbb{R}$.

(i) (*reverse triangle inequality*) $\left|(|x| - |y|)\right| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b| \quad \Rightarrow \quad |a| - |b| \leq |a - b|$$

Switch roles of a and b :

$$|b| - |a| \leq |b - a| = |a - b|$$

(i) follows.

Corollary

Let $x, y \in \mathbb{R}$.

(i) (*reverse triangle inequality*) $\big||x| - |y|\big| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b| \quad \Rightarrow \quad |a| - |b| \leq |a - b|$$

Switch roles of a and b :

$$|b| - |a| \leq |b - a| = |a - b|$$

(i) follows.

(ii): Replace y with $-y$ in standard triangle inequality.



Corollary

Let $x, y \in \mathbb{R}$.

(i) (reverse triangle inequality) $\left| |x| - |y| \right| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b| \quad \Rightarrow \quad |a| - |b| \leq |a - b|$$

Switch roles of a and b :

$$|b| - |a| \leq |b - a| = |a - b|$$

(i) follows.

(ii): Replace y with $-y$ in standard triangle inequality. □

E.g.,

$$2 = 3 - 1 = |3| - |1| \leq |3 - 1| = 2$$

Corollary

Let $x, y \in \mathbb{R}$.

(i) (reverse triangle inequality) $\left|(|x| - |y|)\right| \leq |x - y|$.

(ii) $|x - y| \leq |x| + |y|$.

Proof: (i) Plug $x = a - b$ and $y = b$ into the standard triangle inequality:

$$|a| = |a - b + b| \leq |a - b| + |b| \quad \Rightarrow \quad |a| - |b| \leq |a - b|$$

Switch roles of a and b :

$$|b| - |a| \leq |b - a| = |a - b|$$

(i) follows.

(ii): Replace y with $-y$ in standard triangle inequality. □

E.g.,

$$2 = 3 - 1 = |3| - |1| \leq |3 - 1| = 2$$

$$2 = 3 - 1 = |-3| - |1| \leq |-3 - 1| = 4$$

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \cdots + x_n| \leq |x_1| + |x_2| + \cdots + |x_n|.$$

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof: Induction!

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof: Induction!

Trivial if $n = 1$.

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof: Induction!

Trivial if $n = 1$.

Suppose conclusion holds for n numbers.

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof: Induction!

Trivial if $n = 1$.

Suppose conclusion holds for n numbers.

Consider $n + 1$ numbers x_1, x_2, \dots, x_{n+1} .

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof: Induction!

Trivial if $n = 1$.

Suppose conclusion holds for n numbers.

Consider $n + 1$ numbers x_1, x_2, \dots, x_{n+1} .

$$|x_1 + x_2 + \dots + x_n + x_{n+1}| \leq |x_1 + x_2 + \dots + x_n| + |x_{n+1}|$$

Corollary

Let $x_1, x_2, \dots, x_n \in \mathbb{R}$. Then

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|.$$

Proof: Induction!

Trivial if $n = 1$.

Suppose conclusion holds for n numbers.

Consider $n + 1$ numbers x_1, x_2, \dots, x_{n+1} .

$$\begin{aligned} |x_1 + x_2 + \dots + x_n + x_{n+1}| &\leq |x_1 + x_2 + \dots + x_n| + |x_{n+1}| \\ &\leq |x_1| + |x_2| + \dots + |x_n| + |x_{n+1}| \end{aligned}$$

□

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1|$$

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

$|x|$ is largest when $|x| = 5$.

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

$|x|$ is largest when $|x| = 5$.

So a possible $M = 5^2 + 9(5) + 1 = 71$.

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

$|x|$ is largest when $|x| = 5$.

So a possible $M = 5^2 + 9(5) + 1 = 71$.

Other M work. E.g., $M = 100$ works.

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

$|x|$ is largest when $|x| = 5$.

So a possible $M = 5^2 + 9(5) + 1 = 71$.

Other M work. E.g., $M = 100$ works.

A little calculus shows $M = 77/4 = 19.25$ would also work (least such M).

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

$|x|$ is largest when $|x| = 5$.

So a possible $M = 5^2 + 9(5) + 1 = 71$.

Other M work. E.g., $M = 100$ works.

A little calculus shows $M = 77/4 = 19.25$ would also work (least such M).

We didn't ask for the best M , any M that works will do.

Example: Find a number M such that $|x^2 - 9x + 1| \leq M$ for all $-1 \leq x \leq 5$.

$$|x^2 - 9x + 1| \leq |x^2| + |9x| + |1| = |x|^2 + 9|x| + 1.$$

$|x|^2 + 9|x| + 1$ is largest when $|x|$ is largest (why?).

$|x|$ is largest when $|x| = 5$.

So a possible $M = 5^2 + 9(5) + 1 = 71$.

Other M work. E.g., $M = 100$ works.

A little calculus shows $M = 77/4 = 19.25$ would also work (least such M).

We didn't ask for the best M , any M that works will do.

That sort of point comes up often in analysis.

Definition

$f: D \rightarrow \mathbb{R}$ is *bounded* if $\exists M$ such that $|f(x)| \leq M$ for all $x \in D$.

Definition

$f: D \rightarrow \mathbb{R}$ is *bounded* if $\exists M$ such that $|f(x)| \leq M$ for all $x \in D$.

We proved $x^2 - 9x + 1$ is bounded when the domain is $D = \{x : -1 \leq x \leq 5\}$.

Definition

$f: D \rightarrow \mathbb{R}$ is *bounded* if $\exists M$ such that $|f(x)| \leq M$ for all $x \in D$.

We proved $x^2 - 9x + 1$ is bounded when the domain is $D = \{x : -1 \leq x \leq 5\}$.

If we make $D = \mathbb{R}$, then $x^2 - 9x + 1$ is not bounded.

Definition

$f: D \rightarrow \mathbb{R}$ is *bounded* if $\exists M$ such that $|f(x)| \leq M$ for all $x \in D$.

We proved $x^2 - 9x + 1$ is bounded when the domain is $D = \{x : -1 \leq x \leq 5\}$.

If we make $D = \mathbb{R}$, then $x^2 - 9x + 1$ is not bounded.

We write: $\sup_{x \in D} f(x) := \sup f(D)$ $\inf_{x \in D} f(x) := \inf f(D)$

Definition

$f: D \rightarrow \mathbb{R}$ is *bounded* if $\exists M$ such that $|f(x)| \leq M$ for all $x \in D$.

We proved $x^2 - 9x + 1$ is bounded when the domain is $D = \{x : -1 \leq x \leq 5\}$.

If we make $D = \mathbb{R}$, then $x^2 - 9x + 1$ is not bounded.

We write: $\sup_{x \in D} f(x) := \sup f(D)$ $\inf_{x \in D} f(x) := \inf f(D)$

We also write: $\sup_{-1 \leq x \leq 5} (x^2 - 9x + 1) = 11$ $\inf_{-1 \leq x \leq 5} (x^2 - 9x + 1) = -77/4$.

Definition


$f: D \rightarrow \mathbb{R}$ is *bounded* if $\exists M$ such that $|f(x)| \leq M$ for all $x \in D$.

We proved $x^2 - 9x + 1$ is bounded when the domain is $D = \{x : -1 \leq x \leq 5\}$.

If we make $D = \mathbb{R}$, then $x^2 - 9x + 1$ is not bounded.

We write: $\sup_{x \in D} f(x) := \sup f(D)$ $\inf_{x \in D} f(x) := \inf f(D)$

We also write: $\sup_{-1 \leq x \leq 5} (x^2 - 9x + 1) = 11$ $\inf_{-1 \leq x \leq 5} (x^2 - 9x + 1) = -77/4$.



Proposition

If $f: D \rightarrow \mathbb{R}$, $g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS.

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Suppose b is an upper bound for $g(D)$.

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Suppose b is an upper bound for $g(D)$.

$\Rightarrow f(x) \leq g(x) \leq b$ for all $x \in D$

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Suppose b is an upper bound for $g(D)$.

$\Rightarrow f(x) \leq g(x) \leq b$ for all $x \in D \Rightarrow b$ is an upper bound for $f(D)$.

Proposition

If $f: D \rightarrow \mathbb{R}$, $g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Suppose b is an upper bound for $g(D)$.

$\Rightarrow f(x) \leq g(x) \leq b$ for all $x \in D \Rightarrow b$ is an upper bound for $f(D)$.

Taking the LUB of $g(D)$ gets $f(x) \leq \sup_{y \in D} g(y)$ for any $x \in D$.

Proposition

If $f: D \rightarrow \mathbb{R}$, $g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Suppose b is an upper bound for $g(D)$.

$\Rightarrow f(x) \leq g(x) \leq b$ for all $x \in D \Rightarrow b$ is an upper bound for $f(D)$.

Taking the LUB of $g(D)$ gets $f(x) \leq \sup_{y \in D} g(y)$ for any $x \in D$.

So the LUB of $g(D)$ is an upper bound for $f(D)$.

Proposition

If $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ ($D \neq \emptyset$) are bounded and $f(x) \leq g(x)$ for all $x \in D$, then

$$\sup_{x \in D} f(x) \leq \sup_{x \in D} g(x) \quad \text{and} \quad \inf_{x \in D} f(x) \leq \inf_{x \in D} g(x).$$

Caution: The x on the LHS is different than the x on the RHS. Perhaps think

$$\sup_{x \in D} f(x) \leq \sup_{y \in D} g(y).$$

Proof: We prove the sup, leave the inf as exercise.

Suppose b is an upper bound for $g(D)$.

$\Rightarrow f(x) \leq g(x) \leq b$ for all $x \in D \Rightarrow b$ is an upper bound for $f(D)$.

Taking the LUB of $g(D)$ gets $f(x) \leq \sup_{y \in D} g(y)$ for any $x \in D$.

So the LUB of $g(D)$ is an upper bound for $f(D)$.

$\Rightarrow \sup_{x \in D} f(x) \leq \sup_{y \in D} g(y)$



Caution: $f(x) \leq g(x)$ for all $x \in D$ does **NOT** imply $\sup_{x \in D} f(x) \leq \inf_{y \in D} g(y)$.

Caution: $f(x) \leq g(x)$ for all $x \in D$ does **NOT** imply $\sup_{x \in D} f(x) \leq \inf_{y \in D} g(y)$.

For this you need every value of $g(y)$ to be a bound for f :

Caution: $f(x) \leq g(x)$ for all $x \in D$ does **NOT** imply $\sup_{x \in D} f(x) \leq \inf_{y \in D} g(y)$.

For this you need every value of $g(y)$ to be a bound for f :

That is, $f(x) \leq g(y)$ for all $x, y \in D$ does imply $\sup_{x \in D} f(x) \leq \inf_{y \in D} g(y)$.

Caution: $f(x) \leq g(x)$ for all $x \in D$ does **NOT** imply $\sup_{x \in D} f(x) \leq \inf_{y \in D} g(y)$.

For this you need every value of $g(y)$ to be a bound for f :

That is, $f(x) \leq g(y)$ for all $x, y \in D$ does imply $\sup_{x \in D} f(x) \leq \inf_{y \in D} g(y)$.

Proof is an exercise.

Exercise: Let D be nonempty and $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ be bounded.

Exercise: Let D be nonempty and $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ be bounded.

a) Show

$$\sup_{x \in D} (f(x) + g(x)) \leq \sup_{x \in D} f(x) + \sup_{x \in D} g(x)$$

Exercise: Let D be nonempty and $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ be bounded.

a) Show

$$\sup_{x \in D} (f(x) + g(x)) \leq \sup_{x \in D} f(x) + \sup_{x \in D} g(x)$$

and

$$\inf_{x \in D} (f(x) + g(x)) \geq \inf_{x \in D} f(x) + \inf_{x \in D} g(x).$$

Exercise: Let D be nonempty and $f: D \rightarrow \mathbb{R}, g: D \rightarrow \mathbb{R}$ be bounded.

a) Show

$$\sup_{x \in D} (f(x) + g(x)) \leq \sup_{x \in D} f(x) + \sup_{x \in D} g(x)$$

and

$$\inf_{x \in D} (f(x) + g(x)) \geq \inf_{x \in D} f(x) + \inf_{x \in D} g(x).$$

b) Find examples where we obtain strict inequalities.