

BA: 3.4

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Proof: Suppose $0 \leq x, c \leq 1$. Then

$$|x^2 - c^2|$$

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Proof: Suppose $0 \leq x, c \leq 1$. Then

$$|x^2 - c^2| = |x + c| |x - c|$$

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Proof: Suppose $0 \leq x, c \leq 1$. Then

$$|x^2 - c^2| = |x + c| |x - c| \leq (|x| + |c|) |x - c|$$

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Proof: Suppose $0 \leq x, c \leq 1$. Then

$$|x^2 - c^2| = |x + c| |x - c| \leq (|x| + |c|) |x - c| \leq (1 + 1) |x - c|.$$

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Proof: Suppose $0 \leq x, c \leq 1$. Then

$$|x^2 - c^2| = |x + c| |x - c| \leq (|x| + |c|) |x - c| \leq (1 + 1) |x - c|.$$

Given $\epsilon > 0$, let $\delta := \epsilon/2$.

Definition

Let $S \subset \mathbb{R}$ and $f: S \rightarrow \mathbb{R}$ be a function. Suppose for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $x, c \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$. Then we say f is *uniformly continuous*.

Here δ does not depend on c , only on ϵ .

uniformly continuous \Rightarrow continuous (but not the other way around)

Example: $f: [0, 1] \rightarrow \mathbb{R}$ defined by $f(x) := x^2$ is uniformly continuous.

Proof: Suppose $0 \leq x, c \leq 1$. Then

$$|x^2 - c^2| = |x + c| |x - c| \leq (|x| + |c|) |x - c| \leq (1 + 1) |x - c|.$$

Given $\epsilon > 0$, let $\delta := \epsilon/2$.

If $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.



Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2|$

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.

For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c|$

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.

For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2$

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \ \forall x > 0$,

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$



Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y|$

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|}$

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Suppose $\epsilon < 1$. Consider some $\delta > 0$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Suppose $\epsilon < 1$. Consider some $\delta > 0$.

If $x \in (0, 1)$ and $y = x + \delta/2 \in (0, 1)$, then $|x - y| = \delta/2 < \delta$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Suppose $\epsilon < 1$. Consider some $\delta > 0$.

If $x \in (0, 1)$ and $y = x + \delta/2 \in (0, 1)$, then $|x - y| = \delta/2 < \delta$.

Suppose f is uniformly continuous and δ satisfies the definition.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Suppose $\epsilon < 1$. Consider some $\delta > 0$.

If $x \in (0, 1)$ and $y = x + \delta/2 \in (0, 1)$, then $|x - y| = \delta/2 < \delta$.

Suppose f is uniformly continuous and δ satisfies the definition.

Then $\delta/2 < x(x + \delta/2)\epsilon < x$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \quad \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Suppose $\epsilon < 1$. Consider some $\delta > 0$.

If $x \in (0, 1)$ and $y = x + \delta/2 \in (0, 1)$, then $|x - y| = \delta/2 < \delta$.

Suppose f is uniformly continuous and δ satisfies the definition.

Then $\delta/2 < x(x + \delta/2)\epsilon < x$.

$\delta/2 < x$ for all $x > 0 \quad \Rightarrow \quad \delta \leq 0$.

Example: $g: \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) := x^2$ is not uniformly continuous.

Proof: Suppose it is. Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $|x - c| < \delta$, then $|x^2 - c^2| < \epsilon$.
For $x > 0$ let $c := x + \delta/2$. Then $\epsilon > |x^2 - c^2| = |x + c| |x - c| = (2x + \delta/2)\delta/2 \geq \delta x$.
 $\Rightarrow x < \epsilon/\delta \forall x > 0, \quad \Rightarrow \Leftarrow$

□

Example: $f: (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is not uniformly continuous.

Proof: For any $x, y \in (0, 1)$, $|1/x - 1/y| = \frac{|y-x|}{|xy|} = \frac{|y-x|}{xy}$.

Given $\epsilon > 0$, $\epsilon > |1/x - 1/y| \Leftrightarrow |x - y| < xy\epsilon$.

Suppose $\epsilon < 1$. Consider some $\delta > 0$.

If $x \in (0, 1)$ and $y = x + \delta/2 \in (0, 1)$, then $|x - y| = \delta/2 < \delta$.

Suppose f is uniformly continuous and δ satisfies the definition.

Then $\delta/2 < x(x + \delta/2)\epsilon < x$.

$\delta/2 < x$ for all $x > 0 \quad \Rightarrow \quad \delta \leq 0. \quad \Rightarrow \Leftarrow$

□

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c|$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c|$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c|$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$1/n_k \rightarrow 0$ and $|x_{n_k} - c| \rightarrow 0$ as $k \rightarrow \infty$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \quad \Rightarrow \quad y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \quad \Rightarrow \quad y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$|f(x_{n_k}) - f(c)|$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \quad \Rightarrow \quad y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$|f(x_{n_k}) - f(c)| = |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)|$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \implies y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \end{aligned}$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \quad \Rightarrow \quad y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \geq \epsilon - |f(y_{n_k}) - f(c)| \end{aligned}$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \implies y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \geq \epsilon - |f(y_{n_k}) - f(c)| \end{aligned}$$

$$\text{Or } |f(x_{n_k}) - f(c)| + |f(y_{n_k}) - f(c)| \geq \epsilon.$$

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \implies y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \geq \epsilon - |f(y_{n_k}) - f(c)| \end{aligned}$$

$$\text{Or } |f(x_{n_k}) - f(c)| + |f(y_{n_k}) - f(c)| \geq \epsilon.$$

Either $f(x_{n_k}) \not\rightarrow f(c)$ or $f(y_{n_k}) \not\rightarrow f(c)$, otherwise the LHS would $\rightarrow 0$.

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \Rightarrow y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \geq \epsilon - |f(y_{n_k}) - f(c)| \end{aligned}$$

$$\text{Or } |f(x_{n_k}) - f(c)| + |f(y_{n_k}) - f(c)| \geq \epsilon.$$

Either $f(x_{n_k}) \not\rightarrow f(c)$ or $f(y_{n_k}) \not\rightarrow f(c)$, otherwise the LHS would $\rightarrow 0$.

$\Rightarrow f$ is not continuous at c

Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^{\infty}, \{y_n\}_{n=1}^{\infty}$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^{\infty}$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \Rightarrow y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \geq \epsilon - |f(y_{n_k}) - f(c)| \end{aligned}$$

$$\text{Or } |f(x_{n_k}) - f(c)| + |f(y_{n_k}) - f(c)| \geq \epsilon.$$

Either $f(x_{n_k}) \not\rightarrow f(c)$ or $f(y_{n_k}) \not\rightarrow f(c)$, otherwise the LHS would $\rightarrow 0$.

$\Rightarrow f$ is not continuous at $c \Rightarrow f$ is not continuous.



Theorem

Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then f is uniformly continuous.

Proof: Suppose f is not uniformly continuous.

$\exists \epsilon > 0$ and $\{x_n\}_{n=1}^\infty, \{y_n\}_{n=1}^\infty$ in $[a, b]$ such that $|x_n - y_n| < 1/n$ and $|f(x_n) - f(y_n)| \geq \epsilon$.

By Bolzano–Weierstrass, \exists a convergent subsequence $\{x_{n_k}\}_{k=1}^\infty$.

Let $c := \lim_{k \rightarrow \infty} x_{n_k}$. $a \leq c \leq b$ as $a \leq x_{n_k} \leq b$ for all k ,

$$|y_{n_k} - c| = |y_{n_k} - x_{n_k} + x_{n_k} - c| \leq |y_{n_k} - x_{n_k}| + |x_{n_k} - c| < 1/n_k + |x_{n_k} - c|.$$

$$1/n_k \rightarrow 0 \text{ and } |x_{n_k} - c| \rightarrow 0 \text{ as } k \rightarrow \infty \Rightarrow y_{n_k} \rightarrow c \text{ as } k \rightarrow \infty.$$

$$\begin{aligned} |f(x_{n_k}) - f(c)| &= |f(x_{n_k}) - f(y_{n_k}) + f(y_{n_k}) - f(c)| \\ &\geq |f(x_{n_k}) - f(y_{n_k})| - |f(y_{n_k}) - f(c)| \geq \epsilon - |f(y_{n_k}) - f(c)| \end{aligned}$$

$$\text{Or } |f(x_{n_k}) - f(c)| + |f(y_{n_k}) - f(c)| \geq \epsilon.$$

Either $f(x_{n_k}) \not\rightarrow f(c)$ or $f(y_{n_k}) \not\rightarrow f(c)$, otherwise the LHS would $\rightarrow 0$.

$\Rightarrow f$ is not continuous at $c \Rightarrow f$ is not continuous. □

Remark: See how closed and bounded is important again.

Uniformly continuous functions preserve Cauchy sequences:

Lemma

Let $f: S \rightarrow \mathbb{R}$ be uniformly continuous. Let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence in S . Then $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy.

Uniformly continuous functions preserve Cauchy sequences:

Lemma

Let $f: S \rightarrow \mathbb{R}$ be uniformly continuous. Let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence in S . Then $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy.

Proof: Let $\epsilon > 0$ be given.

Uniformly continuous functions preserve Cauchy sequences:

Lemma

Let $f: S \rightarrow \mathbb{R}$ be uniformly continuous. Let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence in S . Then $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy.

Proof: Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $|f(x) - f(y)| < \epsilon$ whenever $x, y \in S$ and $|x - y| < \delta$.

Uniformly continuous functions preserve Cauchy sequences:

Lemma

Let $f: S \rightarrow \mathbb{R}$ be uniformly continuous. Let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence in S . Then $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy.

Proof: Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $|f(x) - f(y)| < \epsilon$ whenever $x, y \in S$ and $|x - y| < \delta$.

Find $M \in \mathbb{N}$ such that for all $n, k \geq M$, we have $|x_n - x_k| < \delta$.

Uniformly continuous functions preserve Cauchy sequences:

Lemma

Let $f: S \rightarrow \mathbb{R}$ be uniformly continuous. Let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence in S . Then $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy.

Proof: Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $|f(x) - f(y)| < \epsilon$ whenever $x, y \in S$ and $|x - y| < \delta$.

Find $M \in \mathbb{N}$ such that for all $n, k \geq M$, we have $|x_n - x_k| < \delta$.

$\Rightarrow \forall n, k \geq M \quad |f(x_n) - f(x_k)| < \epsilon.$



The lemma gives extension to endpoints for uniformly continuous functions:

Proposition

$f: (a, b) \rightarrow \mathbb{R}$ is uniformly continuous if and only if the limits

$$L_a := \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad L_b := \lim_{x \rightarrow b^-} f(x)$$

exist and $\tilde{f}: [a, b] \rightarrow \mathbb{R}$ defined by

$$\tilde{f}(x) := \begin{cases} f(x) & \text{if } x \in (a, b), \\ L_a & \text{if } x = a, \\ L_b & \text{if } x = b \end{cases}$$

is continuous.

The lemma gives extension to endpoints for uniformly continuous functions:

Proposition

$f: (a, b) \rightarrow \mathbb{R}$ is uniformly continuous if and only if the limits

$$L_a := \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad L_b := \lim_{x \rightarrow b^-} f(x)$$

exist and $\tilde{f}: [a, b] \rightarrow \mathbb{R}$ defined by

$$\tilde{f}(x) := \begin{cases} f(x) & \text{if } x \in (a, b), \\ L_a & \text{if } x = a, \\ L_b & \text{if } x = b \end{cases}$$

is continuous.

Proof: \Leftarrow) If \tilde{f} is continuous, then it is uniformly continuous by the theorem.

The lemma gives extension to endpoints for uniformly continuous functions:

Proposition

$f: (a, b) \rightarrow \mathbb{R}$ is uniformly continuous if and only if the limits

$$L_a := \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad L_b := \lim_{x \rightarrow b^-} f(x)$$

exist and $\tilde{f}: [a, b] \rightarrow \mathbb{R}$ defined by

$$\tilde{f}(x) := \begin{cases} f(x) & \text{if } x \in (a, b), \\ L_a & \text{if } x = a, \\ L_b & \text{if } x = b \end{cases}$$

is continuous.

Proof: \Leftarrow) If \tilde{f} is continuous, then it is uniformly continuous by the theorem.

f is the restriction of \tilde{f} to (a, b) , so f is also uniformly continuous (exercise).

\Rightarrow) Suppose f is uniformly continuous.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.
Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$,

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$,

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n|$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n|$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n|$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2|$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2|$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon.$

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2. \Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$. $\Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists. L_b is almost the same.

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$. $\Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists. L_b is almost the same.

As $\lim_{x \rightarrow a} f(x) = L_a$ exists, $\lim_{x \rightarrow a} \widetilde{f}(x)$ exists and equals L_a (Proposition 3.1.15)

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^{\infty}$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^{\infty}$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^{\infty}$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$. $\Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists. L_b is almost the same.

As $\lim_{x \rightarrow a} f(x) = L_a$ exists, $\lim_{x \rightarrow a} \widetilde{f}(x)$ exists and equals L_a (Proposition 3.1.15) (similarly for b).

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^\infty$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^\infty$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^\infty$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$. $\Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists. L_b is almost the same.

As $\lim_{x \rightarrow a} f(x) = L_a$ exists, $\lim_{x \rightarrow a} \widetilde{f}(x)$ exists and equals L_a (Proposition 3.1.15) (similarly for b).

$\Rightarrow \widetilde{f}$ is continuous at a and b .

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^\infty$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^\infty$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^\infty$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$. $\Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists. L_b is almost the same.

As $\lim_{x \rightarrow a} f(x) = L_a$ exists, $\lim_{x \rightarrow a} \widetilde{f}(x)$ exists and equals L_a (Proposition 3.1.15) (similarly for b).

$\Rightarrow \widetilde{f}$ is continuous at a and b .

As f is continuous at any $c \in (a, b)$, then \widetilde{f} is continuous at $c \in (a, b)$ (Proposition 3.1.15). \square

\Rightarrow) Suppose f is uniformly continuous. WTS that L_a and L_b exist.

Take $\{x_n\}_{n=1}^\infty$ in (a, b) such that $x_n \rightarrow a$ (and so also Cauchy).

Lemma says $\{f(x_n)\}_{n=1}^\infty$ is Cauchy and thus convergent, so let $L_1 := \lim_{n \rightarrow \infty} f(x_n)$.

Take another $\{y_n\}_{n=1}^\infty$ in (a, b) s.t. $y_n \rightarrow a$. Similarly, $L_2 := \lim_{n \rightarrow \infty} f(y_n)$ exists. WTS $L_1 = L_2$.

Given $\epsilon > 0$, find $\delta > 0$ s.t. $|x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon/3$.

$\exists M \in \mathbb{N}$ s.t. for $n \geq M$, $|a - x_n| < \delta/2$, $|a - y_n| < \delta/2$, $|f(x_n) - L_1| < \epsilon/3$, and $|f(y_n) - L_2| < \epsilon/3$.

\Rightarrow for $n \geq M$, $|x_n - y_n| = |x_n - a + a - y_n| \leq |x_n - a| + |a - y_n| < \delta/2 + \delta/2 = \delta$.

$\Rightarrow |L_1 - L_2| = |L_1 - f(x_n) + f(x_n) - f(y_n) + f(y_n) - L_2|$
 $\leq |L_1 - f(x_n)| + |f(x_n) - f(y_n)| + |f(y_n) - L_2| \leq \epsilon/3 + \epsilon/3 + \epsilon/3 = \epsilon$.

$\Rightarrow L_1 = L_2$. $\Rightarrow L_a = \lim_{x \rightarrow a} f(x)$ exists. L_b is almost the same.

As $\lim_{x \rightarrow a} f(x) = L_a$ exists, $\lim_{x \rightarrow a} \widetilde{f}(x)$ exists and equals L_a (Proposition 3.1.15) (similarly for b).

$\Rightarrow \widetilde{f}$ is continuous at a and b .

As f is continuous at any $c \in (a, b)$, then \widetilde{f} is continuous at $c \in (a, b)$ (Proposition 3.1.15). \square

Typical application: if $f: (-1, 0) \cup (0, 1) \rightarrow \mathbb{R}$ is uniformly continuous, then $\lim_{x \rightarrow 0} f(x)$ exists and f has a *removable singularity*: it can be continuously extended to $(-1, 1)$.

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given.

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$,

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$, $|f(x) - f(y)|$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$, $|f(x) - f(y)| \leq K |x - y|$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$, $|f(x) - f(y)| \leq K |x - y| < K\delta$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$, $|f(x) - f(y)| \leq K |x - y| < K\delta = K \frac{\epsilon}{K}$

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$, $|f(x) - f(y)| \leq K |x - y| < K\delta = K \frac{\epsilon}{K} = \epsilon$.

Definition

A function $f: S \rightarrow \mathbb{R}$ is *Lipschitz continuous*, if there exists a $K \in \mathbb{R}$, such that

$$|f(x) - f(y)| \leq K |x - y| \quad \text{for all } x \text{ and } y \text{ in } S.$$

Example: \sin and \cos are Lipschitz with $K = 1$ as

$$|\sin(x) - \sin(y)| \leq |x - y| \quad \text{and} \quad |\cos(x) - \cos(y)| \leq |x - y|.$$

Proposition

A Lipschitz continuous function is uniformly continuous.

Proof: Suppose $f: S \rightarrow \mathbb{R}$ and K are such that

$$|f(x) - f(y)| \leq K |x - y| \text{ for all } x, y \text{ in } S.$$

Let $\epsilon > 0$ be given. Take $\delta := \epsilon/K$.

For all $x, y \in S$ such that $|x - y| < \delta$, $|f(x) - f(y)| \leq K |x - y| < K\delta = K \frac{\epsilon}{K} = \epsilon$.

$\Rightarrow f$ is uniformly continuous.



Geometric interpretation of Lipschitz:

Geometric interpretation of Lipschitz:

Suppose f is Lipschitz with constant K .

Geometric interpretation of Lipschitz:

Suppose f is Lipschitz with constant K .


If $x \neq y$, then $\left| \frac{f(x) - f(y)}{x - y} \right| \leq K$.

Geometric interpretation of Lipschitz:

Suppose f is Lipschitz with constant K .

If $x \neq y$, then $\left| \frac{f(x) - f(y)}{x - y} \right| \leq K$.

$\frac{f(x) - f(y)}{x - y}$ is the slope of the line between $(x, f(x))$ and $(y, f(y))$, the *secant line*.



Geometric interpretation of Lipschitz:

Suppose f is Lipschitz with constant K .


If $x \neq y$, then $\left| \frac{f(x) - f(y)}{x - y} \right| \leq K$.

$\frac{f(x) - f(y)}{x - y}$ is the slope of the line between $(x, f(x))$ and $(y, f(y))$, the *secant line*.

f is Lipschitz

\Leftrightarrow

every secant line has $|\text{slope}| \leq K$.



Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}|$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right|$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}.$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}|$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right|$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \forall x, y \geq 0.$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \quad \forall x, y \geq 0.$

Set $y = 0 \quad \Rightarrow \quad \sqrt{x} \leq Kx$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \quad \forall x, y \geq 0.$

Set $y = 0 \quad \Rightarrow \quad \sqrt{x} \leq Kx \quad \text{in other words, } 1/K \leq \sqrt{x} \quad \text{or} \quad 1/K^2 \leq x.$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \quad \forall x, y \geq 0.$

Set $y = 0 \quad \Rightarrow \quad \sqrt{x} \leq Kx \quad \text{in other words, } 1/K \leq \sqrt{x} \quad \text{or} \quad 1/K^2 \leq x.$

Cannot be true for all $x > 0 \quad \Rightarrow \quad \text{no } K \text{ exists}$

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \quad \Rightarrow \quad \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□

Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \quad \forall x, y \geq 0.$

Set $y = 0 \quad \Rightarrow \quad \sqrt{x} \leq Kx \quad \text{in other words, } 1/K \leq \sqrt{x} \quad \text{or} \quad 1/K^2 \leq x.$

Cannot be true for all $x > 0 \quad \Rightarrow \quad \text{no } K \text{ exists} \quad \Rightarrow \quad g \text{ not Lipschitz.}$

□

Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \Rightarrow \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□


Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \quad \forall x, y \geq 0.$

Set $y = 0 \Rightarrow \sqrt{x} \leq Kx$ in other words, $1/K \leq \sqrt{x}$ or $1/K^2 \leq x.$

Cannot be true for all $x > 0 \Rightarrow$ no K exists $\Rightarrow g$ not Lipschitz.

□



Example: $f: [1, \infty) \rightarrow \mathbb{R}$ given by $f(x) := \sqrt{x}$ is Lipschitz continuous.

Proof: $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| = \frac{|x-y|}{\sqrt{x}+\sqrt{y}}. \quad x \geq 1 \text{ and } y \geq 1 \Rightarrow \frac{1}{\sqrt{x}+\sqrt{y}} \leq \frac{1}{2}.$

Therefore, $|\sqrt{x} - \sqrt{y}| = \left| \frac{x-y}{\sqrt{x}+\sqrt{y}} \right| \leq \frac{1}{2} |x - y|.$

□


Example: $g: [0, \infty) \rightarrow \mathbb{R}$ given by $g(x) := \sqrt{x}$ is **not** Lipschitz continuous.

Proof: Suppose K exists (assume $K > 0$), s.t. $|\sqrt{x} - \sqrt{y}| \leq K |x - y| \quad \forall x, y \geq 0.$

Set $y = 0 \Rightarrow \sqrt{x} \leq Kx$ in other words, $1/K \leq \sqrt{x}$ or $1/K^2 \leq x.$

Cannot be true for all $x > 0 \Rightarrow$ no K exists $\Rightarrow g$ not Lipschitz.

□



Remark: g is uniformly continuous (exercise), but not Lipschitz.