

# BA: 5.2

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Lemma (Additivity)

*Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function.*

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f}$$

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ , and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f)$$

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i$$

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i = \sum_{i=1}^k m_i \Delta x_i + \sum_{i=k+1}^n m_i \Delta x_i$$

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i = \sum_{i=1}^k m_i \Delta x_i + \sum_{i=k+1}^n m_i \Delta x_i = L(P_1, f) + L(P_2, f).$$

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i = \sum_{i=1}^k m_i \Delta x_i + \sum_{i=k+1}^n m_i \Delta x_i = L(P_1, f) + L(P_2, f).$$

sup of RHS over all  $P_1$  and  $P_2$  is like sup of LHS over all  $P$  s.t.  $b \in P$ .

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i = \sum_{i=1}^k m_i \Delta x_i + \sum_{i=k+1}^n m_i \Delta x_i = L(P_1, f) + L(P_2, f).$$

sup of RHS over all  $P_1$  and  $P_2$  is like sup of LHS over all  $P$  s.t.  $b \in P$ .

If  $P = Q \cup \{b\}$ , then  $P$  is a refinement of  $P$ , so  $L(Q, f) \leq L(P, f)$ .

## Lemma (Additivity)

Suppose  $a < b < c$  and  $f: [a, c] \rightarrow \mathbb{R}$  is a bounded function. Then

$$\underline{\int_a^c} f = \underline{\int_a^b} f + \underline{\int_b^c} f \quad \text{and} \quad \overline{\int_a^c} f = \overline{\int_a^b} f + \overline{\int_b^c} f.$$

**Proof:** Let  $P_1 = \{x_0, x_1, \dots, x_k\}$  be a partition of  $[a, b]$ ,  
and  $P_2 = \{x_k, x_{k+1}, \dots, x_n\}$  of  $[b, c]$ ,  
 $\Rightarrow P := P_1 \cup P_2 = \{x_0, x_1, \dots, x_n\}$  is a partition of  $[a, c]$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i = \sum_{i=1}^k m_i \Delta x_i + \sum_{i=k+1}^n m_i \Delta x_i = L(P_1, f) + L(P_2, f).$$

sup of RHS over all  $P_1$  and  $P_2$  is like sup of LHS over all  $P$  s.t.  $b \in P$ .

If  $P = Q \cup \{b\}$ , then  $P$  is a refinement of  $Q$ , so  $L(Q, f) \leq L(P, f)$ .

$\Rightarrow$  sup  $L(P, f)$  over all  $P$  with  $b \in P$  is sufficient to get sup  $L(P, f)$  over all  $P$ .

$$\underline{\int_a^c f}$$

$$\underline{\int_a^c} f = \sup \{L(P, f) : P \text{ a partition of } [a, c]\}$$

$$\underline{\int_a^c f} = \sup \{L(P, f) : P \text{ a partition of } [a, c]\}$$
$$= \sup \{L(P, f) : P \text{ a partition of } [a, c], b \in P\}$$

$$\underline{\int_a^c} f = \sup \{L(P, f) : P \text{ a partition of } [a, c]\}$$

$$= \sup \{L(P, f) : P \text{ a partition of } [a, c], b \in P\}$$

$$= \sup \{L(P_1, f) + L(P_2, f) : P_1 \text{ a partition of } [a, b], P_2 \text{ a partition of } [b, c]\}$$

$$\begin{aligned}
\underline{\int_a^c} f &= \sup \{L(P, f) : P \text{ a partition of } [a, c]\} \\
&= \sup \{L(P, f) : P \text{ a partition of } [a, c], b \in P\} \\
&= \sup \{L(P_1, f) + L(P_2, f) : P_1 \text{ a partition of } [a, b], P_2 \text{ a partition of } [b, c]\} \\
&= \sup \{L(P_1, f) : P_1 \text{ a partition of } [a, b]\} \\
&\quad + \sup \{L(P_2, f) : P_2 \text{ a partition of } [b, c]\}
\end{aligned}$$

$$\begin{aligned}
\underline{\int_a^c} f &= \sup \{L(P, f) : P \text{ a partition of } [a, c]\} \\
&= \sup \{L(P, f) : P \text{ a partition of } [a, c], b \in P\} \\
&= \sup \{L(P_1, f) + L(P_2, f) : P_1 \text{ a partition of } [a, b], P_2 \text{ a partition of } [b, c]\} \\
&= \sup \{L(P_1, f) : P_1 \text{ a partition of } [a, b]\} \\
&\quad + \sup \{L(P_2, f) : P_2 \text{ a partition of } [b, c]\} \\
&= \underline{\int_a^b} f + \underline{\int_b^c} f.
\end{aligned}$$

$$\begin{aligned}
\underline{\int_a^c} f &= \sup \{L(P, f) : P \text{ a partition of } [a, c]\} \\
&= \sup \{L(P, f) : P \text{ a partition of } [a, c], b \in P\} \\
&= \sup \{L(P_1, f) + L(P_2, f) : P_1 \text{ a partition of } [a, b], P_2 \text{ a partition of } [b, c]\} \\
&= \sup \{L(P_1, f) : P_1 \text{ a partition of } [a, b]\} \\
&\quad + \sup \{L(P_2, f) : P_2 \text{ a partition of } [b, c]\} \\
&= \underline{\int_a^b} f + \underline{\int_b^c} f.
\end{aligned}$$

The upper integral argument is analogous (see book).



## Proposition (Additivity)

*Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ .*

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f = \underline{\int_a^c f}$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f}$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f}$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f}$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f} = \int_a^c f$ .

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

The lemma  $\Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f} = \int_a^c f.$

$$\Rightarrow \underline{\int_a^b f} + \underline{\int_b^c f} = \overline{\int_a^b f} + \overline{\int_b^c f}.$$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

$$\text{The lemma} \Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f} = \int_a^c f.$$

$$\Rightarrow \underline{\int_a^b f} + \underline{\int_b^c f} = \overline{\int_a^b f} + \overline{\int_b^c f}.$$

$$\underline{\int_a^b f} \leq \overline{\int_a^b f} \text{ and } \underline{\int_b^c f} \leq \overline{\int_b^c f}$$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

$$\text{The lemma} \Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f} = \int_a^c f.$$

$$\Rightarrow \underline{\int_a^b f} + \underline{\int_b^c f} = \overline{\int_a^b f} + \overline{\int_b^c f}.$$

$$\underline{\int_a^b f} \leq \overline{\int_a^b f} \text{ and } \underline{\int_b^c f} \leq \overline{\int_b^c f} \Rightarrow \underline{\int_a^b f} = \overline{\int_a^b f} \text{ and } \underline{\int_b^c f} = \overline{\int_b^c f}.$$

## Proposition (Additivity)

Let  $a < b < c$ .  $f: [a, c] \rightarrow \mathbb{R}$  is Riemann integrable if and only if  $f$  is Riemann integrable on  $[a, b]$  and  $[b, c]$ . If  $f$  is Riemann integrable, then

$$\int_a^c f = \int_a^b f + \int_b^c f.$$

**Proof:** Suppose  $f \in \mathcal{R}[a, c]$ , then  $\overline{\int_a^c f} = \underline{\int_a^c f} = \int_a^c f$ .

$$\text{The lemma} \Rightarrow \int_a^c f = \underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} \leq \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f} = \int_a^c f.$$

$$\Rightarrow \underline{\int_a^b f} + \underline{\int_b^c f} = \overline{\int_a^b f} + \overline{\int_b^c f}.$$

$$\underline{\int_a^b f} \leq \overline{\int_a^b f} \text{ and } \underline{\int_b^c f} \leq \overline{\int_b^c f} \Rightarrow \underline{\int_a^b f} = \overline{\int_a^b f} \text{ and } \underline{\int_b^c f} = \overline{\int_b^c f}.$$

$$\Rightarrow f \text{ is Riemann integrable on } [a, b] \text{ and } [b, c] \text{ and } \int_a^c f = \int_a^b f + \int_b^c f.$$

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f}$$

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f}$$

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} = \int_a^b f + \int_b^c f$$

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} = \int_a^b f + \int_b^c f = \overline{\int_a^b f} + \overline{\int_b^c f}$$

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} = \int_a^b f + \int_b^c f = \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f}.$$

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} = \int_a^b f + \int_b^c f = \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f}.$$

$\Rightarrow f$  is Riemann integrable on  $[a, c]$  and  $\int_a^c f = \int_a^b f + \int_b^c f$ .



Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\underline{\int_a^c f} = \underline{\int_a^b f} + \underline{\int_b^c f} = \int_a^b f + \int_b^c f = \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f}.$$

$\Rightarrow f$  is Riemann integrable on  $[a, c]$  and  $\int_a^c f = \int_a^b f + \int_b^c f$ . □

### Corollary

*If  $f \in \mathcal{R}[a, b]$  and  $[c, d] \subset [a, b]$ , then the restriction  $f|_{[c, d]}$  is in  $\mathcal{R}[c, d]$ .*

Now assume  $f$  is Riemann integrable on  $[a, b]$  and on  $[b, c]$ .

$$\int_a^c f = \int_a^b f + \int_b^c f = \int_a^b f + \int_b^c f = \overline{\int_a^b f} + \overline{\int_b^c f} = \overline{\int_a^c f}.$$

$\Rightarrow f$  is Riemann integrable on  $[a, c]$  and  $\int_a^c f = \int_a^b f + \int_b^c f$ . □

### Corollary

If  $f \in \mathcal{R}[a, b]$  and  $[c, d] \subset [a, b]$ , then the restriction  $f|_{[c, d]}$  is in  $\mathcal{R}[c, d]$ .

**Proof:** Exercise.

## Proposition (Linearity)

*Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .*

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ .

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\}$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\}$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$L(P, \alpha f)$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\underline{\int_a^b \alpha f(x) dx}$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\underline{\int_a^b \alpha f(x) dx} = \sup \{L(P, \alpha f) : P \text{ a partition of } [a, b]\}$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\begin{aligned} \int_a^b \alpha f(x) dx &= \sup \{L(P, \alpha f) : P \text{ a partition of } [a, b]\} \\ &= \sup \{\alpha L(P, f) : P \text{ a partition of } [a, b]\} \end{aligned}$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\begin{aligned} \underline{\int_a^b \alpha f(x) dx} &= \sup \{L(P, \alpha f) : P \text{ a partition of } [a, b]\} \\ &= \sup \{\alpha L(P, f) : P \text{ a partition of } [a, b]\} \\ &= \alpha \sup \{L(P, f) : P \text{ a partition of } [a, b]\} \end{aligned}$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\begin{aligned} \underline{\int_a^b \alpha f(x) dx} &= \sup \{L(P, \alpha f) : P \text{ a partition of } [a, b]\} \\ &= \sup \{\alpha L(P, f) : P \text{ a partition of } [a, b]\} \\ &= \alpha \sup \{L(P, f) : P \text{ a partition of } [a, b]\} = \alpha \underline{\int_a^b f(x) dx}. \end{aligned}$$

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\begin{aligned} \int_a^b \alpha f(x) dx &= \sup \{L(P, \alpha f) : P \text{ a partition of } [a, b]\} \\ &= \sup \{\alpha L(P, f) : P \text{ a partition of } [a, b]\} \\ &= \alpha \sup \{L(P, f) : P \text{ a partition of } [a, b]\} = \alpha \int_a^b f(x) dx. \end{aligned}$$

Similarly,  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

## Proposition (Linearity)

Let  $f$  and  $g$  be in  $\mathcal{R}[a, b]$  and  $\alpha \in \mathbb{R}$ .

(i)  $\alpha f$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b \alpha f(x) dx = \alpha \int_a^b f(x) dx$ .

(ii)  $f + g$  is in  $\mathcal{R}[a, b]$  and  $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$ .

**Proof:** (i) Suppose  $\alpha \geq 0$ . Let  $P$  be a partition of  $[a, b]$ ,  $m_i$  as usual.  
 $\inf\{\alpha f(x) : x \in [x_{i-1}, x_i]\} = \alpha \inf\{f(x) : x \in [x_{i-1}, x_i]\} = \alpha m_i$ .

$$L(P, \alpha f) = \sum_{i=1}^n \alpha m_i \Delta x_i = \alpha \sum_{i=1}^n m_i \Delta x_i = \alpha L(P, f).$$

Similarly,  $U(P, \alpha f) = \alpha U(P, f)$ .

$$\begin{aligned} \underline{\int_a^b \alpha f(x) dx} &= \sup \{L(P, \alpha f) : P \text{ a partition of } [a, b]\} \\ &= \sup \{\alpha L(P, f) : P \text{ a partition of } [a, b]\} \\ &= \alpha \sup \{L(P, f) : P \text{ a partition of } [a, b]\} = \alpha \underline{\int_a^b f(x) dx}. \end{aligned}$$

Similarly,  $\overline{\int_a^b \alpha f(x) dx} = \alpha \overline{\int_a^b f(x) dx} \Rightarrow$  (i) follows for  $\alpha \geq 0$ .

**Exercise:** Finish proof of (i) ( $\alpha < 0$ ):

**Exercise:** Finish proof of (i) ( $\alpha < 0$ ):

Show  $-f$  is Riemann integrable and  $\int_a^b -f(x) dx = -\int_a^b f(x) dx$ .

**Exercise:** Finish proof of (i) ( $\alpha < 0$ ):

Show  $-f$  is Riemann integrable and  $\int_a^b -f(x) dx = -\int_a^b f(x) dx$ .

**Exercise:** Prove (ii).



**Exercise:** Finish proof of (i) ( $\alpha < 0$ ):

Show  $-f$  is Riemann integrable and  $\int_a^b -f(x) dx = -\int_a^b f(x) dx$ .

**Exercise:** Prove (ii).



(ii) is not as trivial as may seem,

**Exercise:** Finish proof of (i) ( $\alpha < 0$ ):

Show  $-f$  is Riemann integrable and  $\int_a^b -f(x) dx = -\int_a^b f(x) dx$ .

**Exercise:** Prove (ii).



(ii) is not as trivial as may seem, see

## Proposition

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded functions. Then

$$\overline{\int_a^b (f+g)} \leq \overline{\int_a^b f} + \overline{\int_a^b g}, \quad \text{and} \quad \underline{\int_a^b (f+g)} \geq \underline{\int_a^b f} + \underline{\int_a^b g}.$$

**Exercise:** Finish proof of (i) ( $\alpha < 0$ ):

Show  $-f$  is Riemann integrable and  $\int_a^b -f(x) dx = -\int_a^b f(x) dx$ .

**Exercise:** Prove (ii).



(ii) is not as trivial as may seem, see

### Proposition

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded functions. Then

$$\overline{\int_a^b (f + g)} \leq \overline{\int_a^b f} + \overline{\int_a^b g}, \quad \text{and} \quad \underline{\int_a^b (f + g)} \geq \underline{\int_a^b f} + \underline{\int_a^b g}.$$

**Proof:** Exercise.

## Proposition (Monotonicity)

*Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ .*

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$L(P, f)$

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i$$

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \tilde{m}_i \Delta x_i$$

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \tilde{m}_i \Delta x_i = L(P, g)$$

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \tilde{m}_i \Delta x_i = L(P, g) \quad \Rightarrow \quad \underline{\int_a^b f} \leq \underline{\int_a^b g}.$$

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \tilde{m}_i \Delta x_i = L(P, g) \quad \Rightarrow \quad \underline{\int_a^b f} \leq \underline{\int_a^b g}.$$

Similarly,  $\overline{\int_a^b f} \leq \overline{\int_a^b g}$ .

## Proposition (Monotonicity)

Let  $f: [a, b] \rightarrow \mathbb{R}$  and  $g: [a, b] \rightarrow \mathbb{R}$  be bounded, and  $f(x) \leq g(x) \forall x \in [a, b]$ . Then

$$\underline{\int_a^b f} \leq \underline{\int_a^b g} \quad \text{and} \quad \overline{\int_a^b f} \leq \overline{\int_a^b g}.$$

Moreover, if  $f$  and  $g$  are in  $\mathcal{R}[a, b]$ , then  $\int_a^b f \leq \int_a^b g$ .

**Proof:** Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition of  $[a, b]$ .

Let  $m_i := \inf \{f(x) : x \in [x_{i-1}, x_i]\}$  and  $\tilde{m}_i := \inf \{g(x) : x \in [x_{i-1}, x_i]\}$ .

$f(x) \leq g(x)$  for all  $x \Rightarrow m_i \leq \tilde{m}_i$ .

$$L(P, f) = \sum_{i=1}^n m_i \Delta x_i \leq \sum_{i=1}^n \tilde{m}_i \Delta x_i = L(P, g) \quad \Rightarrow \quad \underline{\int_a^b f} \leq \underline{\int_a^b g}.$$

Similarly,  $\overline{\int_a^b f} \leq \overline{\int_a^b g}$ .

If  $f$  and  $g$  are integrable,  $\Rightarrow \int_a^b f \leq \int_a^b g$ .



## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.  
Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta,$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)|$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad f(x) - f(y) \leq |f(x) - f(y)|$$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad f(x) - f(y) \leq |f(x) - f(y)| < \frac{\epsilon}{b-a}.$$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad f(x) - f(y) \leq |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$\Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f}$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad f(x) - f(y) \leq |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$\Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} \leq U(P, f) - L(P, f)$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} \leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right)$$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\begin{aligned} \Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} &\leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right) \\ &= \sum_{i=1}^n (M_i - m_i) \Delta x_i \end{aligned}$$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad f(x) - f(y) \leq |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\begin{aligned} \Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} &\leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right) \\ &= \sum_{i=1}^n (M_i - m_i) \Delta x_i < \frac{\epsilon}{b-a} \sum_{i=1}^n \Delta x_i \end{aligned}$$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\begin{aligned} \Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} &\leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right) \\ &= \sum_{i=1}^n (M_i - m_i) \Delta x_i < \frac{\epsilon}{b-a} \sum_{i=1}^n \Delta x_i = \frac{\epsilon}{b-a} (b-a) = \epsilon. \end{aligned}$$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow$   $f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\begin{aligned} \Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} &\leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right) \\ &= \sum_{i=1}^n (M_i - m_i) \Delta x_i < \frac{\epsilon}{b-a} \sum_{i=1}^n \Delta x_i = \frac{\epsilon}{b-a} (b-a) = \epsilon. \end{aligned}$$

$\epsilon > 0$  was arbitrary

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\begin{aligned} \Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} &\leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right) \\ &= \sum_{i=1}^n (M_i - m_i) \Delta x_i < \frac{\epsilon}{b-a} \sum_{i=1}^n \Delta x_i = \frac{\epsilon}{b-a} (b-a) = \epsilon. \end{aligned}$$

$\epsilon > 0$  was arbitrary  $\Rightarrow \overline{\int_a^b f} = \underline{\int_a^b f}$

## Lemma

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f \in \mathcal{R}[a, b]$ .

**Proof:**  $[a, b]$  is closed & bounded  $\Rightarrow f$  is bounded & uniformly continuous.

Given  $\epsilon > 0$ , find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies  $|f(x) - f(y)| < \frac{\epsilon}{b-a}$ .

Let  $P = \{x_0, x_1, \dots, x_n\}$  be a partition s.t.  $\Delta x_i < \delta$  for all  $i$ .

$\forall x, y \in [x_{i-1}, x_i], \quad |x - y| \leq \Delta x_i < \delta, \quad \Rightarrow \quad |f(x) - f(y)| < \frac{\epsilon}{b-a}.$

$f$  continuous so  $f$  attains a maximum on  $[x_{i-1}, x_i]$  at  $x$  and a minimum at  $y$ .

$\Rightarrow f(x) = M_i$  and  $f(y) = m_i$  (usual notation).

$\Rightarrow M_i - m_i = f(x) - f(y) < \frac{\epsilon}{b-a}.$

$$\begin{aligned} \Rightarrow \overline{\int_a^b f} - \underline{\int_a^b f} &\leq U(P, f) - L(P, f) = \left( \sum_{i=1}^n M_i \Delta x_i \right) - \left( \sum_{i=1}^n m_i \Delta x_i \right) \\ &= \sum_{i=1}^n (M_i - m_i) \Delta x_i < \frac{\epsilon}{b-a} \sum_{i=1}^n \Delta x_i = \frac{\epsilon}{b-a} (b-a) = \epsilon. \end{aligned}$$

$\epsilon > 0$  was arbitrary  $\Rightarrow \overline{\int_a^b f} = \underline{\int_a^b f} \Rightarrow f$  is Riemann integrable.

□

It is not hard to prove that functions with finitely many discontinuities are also integrable (we skip the proof),

It is not hard to prove that functions with finitely many discontinuities are also integrable (we skip the proof), although a lot more is true.

It is not hard to prove that functions with finitely many discontinuities are also integrable (we skip the proof), although a lot more is true.

Also,

### Proposition

*Let  $f: [a, b] \rightarrow \mathbb{R}$  be a monotone function. Then  $f \in \mathcal{R}[a, b]$ .*

It is not hard to prove that functions with finitely many discontinuities are also integrable (we skip the proof), although a lot more is true.

Also,

### Proposition

*Let  $f: [a, b] \rightarrow \mathbb{R}$  be a monotone function. Then  $f \in \mathcal{R}[a, b]$ .*

**Proof:** Exercise.

It is not hard to prove that functions with finitely many discontinuities are also integrable (we skip the proof), although a lot more is true.

Also,

### Proposition

*Let  $f: [a, b] \rightarrow \mathbb{R}$  be a monotone function. Then  $f \in \mathcal{R}[a, b]$ .*

**Proof:** Exercise.

Functions  $h$  of the form  $h = f - g$  where  $f$  and  $g$  are monotone are said to be of *bounded variation*.

It is not hard to prove that functions with finitely many discontinuities are also integrable (we skip the proof), although a lot more is true.

Also,

### Proposition

Let  $f: [a, b] \rightarrow \mathbb{R}$  be a monotone function. Then  $f \in \mathcal{R}[a, b]$ .

**Proof:** Exercise.

Functions  $h$  of the form  $h = f - g$  where  $f$  and  $g$  are monotone are said to be of *bounded variation*.

Such functions are Riemann integrable.

**Exercise:** Prove the *mean value theorem for integrals*: If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then there exists a  $c \in [a, b]$  such that  $\int_a^b f = f(c)(b - a)$ .

**Exercise:** Prove the *mean value theorem for integrals*: If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then there exists a  $c \in [a, b]$  such that  $\int_a^b f = f(c)(b - a)$ .

**Exercise:** Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function such that  $f(x) \geq 0$  for all  $x \in [a, b]$  and  $\int_a^b f = 0$ . Prove that  $f(x) = 0$  for all  $x$ .

**Exercise:** Prove the *mean value theorem for integrals*: If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then there exists a  $c \in [a, b]$  such that  $\int_a^b f = f(c)(b - a)$ .

**Exercise:** Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function such that  $f(x) \geq 0$  for all  $x \in [a, b]$  and  $\int_a^b f = 0$ . Prove that  $f(x) = 0$  for all  $x$ .

**Exercise (Challenging):** Suppose  $f \in \mathcal{R}[a, b]$ , then the function that takes  $x$  to  $|f(x)|$  is also Riemann integrable on  $[a, b]$ .

**Exercise:** Prove the *mean value theorem for integrals*: If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous, then there exists a  $c \in [a, b]$  such that  $\int_a^b f = f(c)(b - a)$ .

**Exercise:** Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function such that  $f(x) \geq 0$  for all  $x \in [a, b]$  and  $\int_a^b f = 0$ . Prove that  $f(x) = 0$  for all  $x$ .

**Exercise (Challenging):** Suppose  $f \in \mathcal{R}[a, b]$ , then the function that takes  $x$  to  $|f(x)|$  is also Riemann integrable on  $[a, b]$ . Then show

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$