

BA: 7.4

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy*

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Given $\epsilon > 0$, $\exists M$ such that $\forall n \geq M$, $d(p, x_n) < \epsilon/2$.

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Given $\epsilon > 0$, $\exists M$ such that $\forall n \geq M$, $d(p, x_n) < \epsilon/2$.

$\Rightarrow \forall n, k \geq M, d(x_n, x_k)$

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Given $\epsilon > 0$, $\exists M$ such that $\forall n \geq M$, $d(p, x_n) < \epsilon/2$.

$\Rightarrow \forall n, k \geq M, \quad d(x_n, x_k) \leq d(x_n, p) + d(p, x_k)$

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Given $\epsilon > 0$, $\exists M$ such that $\forall n \geq M$, $d(p, x_n) < \epsilon/2$.

$\Rightarrow \forall n, k \geq M$, $d(x_n, x_k) \leq d(x_n, p) + d(p, x_k) < \epsilon/2 + \epsilon/2$

Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Given $\epsilon > 0$, $\exists M$ such that $\forall n \geq M$, $d(p, x_n) < \epsilon/2$.

$\Rightarrow \forall n, k \geq M$, $d(x_n, x_k) \leq d(x_n, p) + d(p, x_k) < \epsilon/2 + \epsilon/2 = \epsilon$.



Definition

A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space (X, d) is *Cauchy* if for every $\epsilon > 0$, there exists an $M \in \mathbb{N}$ such that for all $n \geq M$ and all $k \geq M$, $d(x_n, x_k) < \epsilon$.

Proposition

A convergent sequence in a metric space is Cauchy.

Proof: Suppose $\{x_n\}_{n=1}^{\infty}$ converges to p .

Given $\epsilon > 0$, $\exists M$ such that $\forall n \geq M$, $d(p, x_n) < \epsilon/2$.

$\Rightarrow \forall n, k \geq M$, $d(x_n, x_k) \leq d(x_n, p) + d(p, x_k) < \epsilon/2 + \epsilon/2 = \epsilon$. □

Definition

A metric space (X, d) is *complete* or *Cauchy-complete* if every Cauchy sequence $\{x_n\}_{n=1}^{\infty}$ in X converges to a $p \in X$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy,

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}|$$

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2}$$

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^\infty$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2}$$

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j)$$

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^{\infty}$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^{\infty}$ is Cauchy.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^\infty$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ is Cauchy.

\mathbb{R} is complete

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^\infty$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ is Cauchy.

\mathbb{R} is complete $\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ converges to some $y_k \in \mathbb{R}$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^\infty$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ is Cauchy.

\mathbb{R} is complete $\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ converges to some $y_k \in \mathbb{R}$.

Write $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$.

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^\infty$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ is Cauchy.

\mathbb{R} is complete $\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ converges to some $y_k \in \mathbb{R}$.

Write $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$.

By a proposition, $\{x_m\}_{m=1}^\infty$ converges to y

Proposition

The space \mathbb{R}^n with the standard metric is a complete metric space.

Already proved for $\mathbb{R} = \mathbb{R}^1$.

Proof: Let $\{x_m\}_{m=1}^\infty$ in \mathbb{R}^n be Cauchy, write $x_m = (x_{m,1}, x_{m,2}, \dots, x_{m,n})$.

Given $\epsilon > 0$, $\exists M$ such that $\forall i, j \geq M$, $d(x_i, x_j) < \epsilon$.

Fix $k = 1, 2, \dots, n$. For $i, j \geq M$,

$$|x_{i,k} - x_{j,k}| = \sqrt{(x_{i,k} - x_{j,k})^2} \leq \sqrt{\sum_{\ell=1}^n (x_{i,\ell} - x_{j,\ell})^2} = d(x_i, x_j) < \epsilon.$$

$\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ is Cauchy.

\mathbb{R} is complete $\Rightarrow \{x_{m,k}\}_{m=1}^\infty$ converges to some $y_k \in \mathbb{R}$.

Write $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$.

By a proposition, $\{x_m\}_{m=1}^\infty$ converges to $y \Rightarrow \mathbb{R}^n$ is complete.



Consider $C([a, b], \mathbb{R})$ with the uniform norm as the metric

$$d(f, g) = \|f - g\|_{[a, b]} = \sup_{x \in [a, b]} |f(x) - g(x)|.$$

Consider $C([a, b], \mathbb{R})$ with the uniform norm as the metric

$$d(f, g) = \|f - g\|_{[a, b]} = \sup_{x \in [a, b]} |f(x) - g(x)|.$$

Convergence in this metric space is uniform convergence.

Consider $C([a, b], \mathbb{R})$ with the uniform norm as the metric

$$d(f, g) = \|f - g\|_{[a, b]} = \sup_{x \in [a, b]} |f(x) - g(x)|.$$

Convergence in this metric space is uniform convergence.

In chapter 6, we proved a uniform limit of continuous functions is continuous.

Consider $C([a, b], \mathbb{R})$ with the uniform norm as the metric

$$d(f, g) = \|f - g\|_{[a, b]} = \sup_{x \in [a, b]} |f(x) - g(x)|.$$

Convergence in this metric space is uniform convergence.

In chapter 6, we proved a uniform limit of continuous functions is continuous.

The following proposition is then an easy exercise in verifying the definitions:

Proposition

The space of continuous functions $C([a, b], \mathbb{R})$ with the uniform norm as metric is a complete metric space.

Example: A subset of a complete metric space need not be complete:

Example: A subset of a complete metric space need not be complete:
 $(0, 1]$ with the subspace metric is not complete.

Example: A subset of a complete metric space need not be complete:
 $(0, 1]$ with the subspace metric is not complete.
 $\{1/n\}_{n=1}^{\infty}$ is Cauchy with no limit in $(0, 1]$.

Example: A subset of a complete metric space need not be complete:
 $(0, 1]$ with the subspace metric is not complete.
 $\{1/n\}_{n=1}^{\infty}$ is Cauchy with no limit in $(0, 1]$.

Proposition

Suppose (X, d) is a complete metric space and $E \subset X$ is closed.

Example: A subset of a complete metric space need not be complete:
 $(0, 1]$ with the subspace metric is not complete.
 $\{1/n\}_{n=1}^{\infty}$ is Cauchy with no limit in $(0, 1]$.

Proposition

Suppose (X, d) is a complete metric space and $E \subset X$ is closed. Then E is a complete metric space with the subspace metric.

Example: A subset of a complete metric space need not be complete:
 $(0, 1]$ with the subspace metric is not complete.
 $\{1/n\}_{n=1}^{\infty}$ is Cauchy with no limit in $(0, 1]$.

Proposition

Suppose (X, d) is a complete metric space and $E \subset X$ is closed. Then E is a complete metric space with the subspace metric.

Proof: Exercise.

Definition

Let (X, d) be a metric space and $K \subset X$.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m}$

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Again, if \exists finite subcover then $(0, 1) \subset U_j$ for some j ,

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Again, if \exists finite subcover then $(0, 1) \subset U_j$ for some j , $\Rightarrow \Leftarrow$

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Again, if \exists finite subcover then $(0, 1) \subset U_j$ for some j , $\Rightarrow \Leftarrow$

Example: $\{0\} \subset \mathbb{R}$ is compact

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Again, if \exists finite subcover then $(0, 1) \subset U_j$ for some j , $\Rightarrow \Leftarrow$

Example: $\{0\} \subset \mathbb{R}$ is compact

Proof: For an open cover $\{U_\lambda\}_{\lambda \in I}$,

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Again, if \exists finite subcover then $(0, 1) \subset U_j$ for some j , $\Rightarrow \Leftarrow$

Example: $\{0\} \subset \mathbb{R}$ is compact

Proof: For an open cover $\{U_\lambda\}_{\lambda \in I}$, $\exists \lambda_0$ s.t. $0 \in U_{\lambda_0}$.

Definition

Let (X, d) be a metric space and $K \subset X$. The set K is *compact* if for every collection of open sets $\{U_\lambda\}_{\lambda \in I}$ such that $K \subset \bigcup_{\lambda \in I} U_\lambda$,

there exists a finite subset $\{\lambda_1, \lambda_2, \dots, \lambda_m\} \subset I$ such that $K \subset \bigcup_{j=1}^m U_{\lambda_j}$.

Call $\{U_\lambda\}_{\lambda \in I}$ an *open cover* of K .

K is compact if *every open cover of K has a finite subcover*.

Example: \mathbb{R} is not compact.

Proof: Take $U_j := (-j, j)$, $j \in \mathbb{N}$. $\bigcup_j (-j, j) = \mathbb{R}$ by Archimedean property.

If $\mathbb{R} \subset U_{j_1} \cup U_{j_2} \cup \dots \cup U_{j_m}$ ($j_1 < j_2 < \dots < j_m$) $\Rightarrow \mathbb{R} \subset U_{j_m} \Rightarrow \Leftarrow$

Example: $(0, 1) \subset \mathbb{R}$ is not compact.

Proof: Consider $U_j := (1/j, 1 - 1/j)$ for $j = 3, 4, 5, \dots$ $(0, 1) = \bigcup_{j=3}^{\infty} U_j$.

Again, if \exists finite subcover then $(0, 1) \subset U_j$ for some j , $\Rightarrow \Leftarrow$

Example: $\{0\} \subset \mathbb{R}$ is compact

Proof: For an open cover $\{U_\lambda\}_{\lambda \in I}$, $\exists \lambda_0$ s.t. $0 \in U_{\lambda_0}$. $\{U_{\lambda_0}\}$ is a finite subcover.

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.

Proof: Fix $p \in X$.

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.

Proof: Fix $p \in X$.

The open cover $K \subset \bigcup_{n=1}^{\infty} B(p, n) = X$

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.

Proof: Fix $p \in X$.

The open cover $K \subset \bigcup_{n=1}^{\infty} B(p, n) = X$

has a finite subcover

$$K \subset \bigcup_{j=1}^m B(p, n_j)$$

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.

Proof: Fix $p \in X$.

The open cover $K \subset \bigcup_{n=1}^{\infty} B(p, n) = X$

has a finite subcover

$$K \subset \bigcup_{j=1}^m B(p, n_j) = B(p, n_m).$$

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.

Proof: Fix $p \in X$.

The open cover $K \subset \bigcup_{n=1}^{\infty} B(p, n) = X$

has a finite subcover

$$K \subset \bigcup_{j=1}^m B(p, n_j) = B(p, n_m).$$

$\Rightarrow K$ is bounded.

Proposition

Let (X, d) be a metric space. If $K \subset X$ is compact, then K is closed and bounded.


Proof: Fix $p \in X$.

The open cover $K \subset \bigcup_{n=1}^{\infty} B(p, n) = X$

has a finite subcover

$$K \subset \bigcup_{j=1}^m B(p, n_j) = B(p, n_m).$$

$\Rightarrow K$ is bounded.



Suppose K is not closed: $\bar{K} \neq K$

Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$

Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n


$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$

Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$




Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$

That's an open cover as $C(x, 1/n)^c$ are open.



Suppose K is not closed: $\bar{K} \neq K$


$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$

That's an open cover as $C(x, 1/n)^c$ are open.

Take a finite collection $\bigcup_{j=1}^m C(x, 1/n_j)^c$



Suppose K is not closed: $\bar{K} \neq K$


$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

$$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$$

That's an open cover as $C(x, 1/n)^c$ are open.

Take a finite collection $\bigcup_{j=1}^m C(x, 1/n_j)^c = C(x, 1/n_m)^c$



Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$


$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

$$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$$

That's an open cover as $C(x, 1/n)^c$ are open.

Take a finite collection $\bigcup_{j=1}^m C(x, 1/n_j)^c = C(x, 1/n_m)^c$

$x \in \bar{K}$



Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K$.


$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n

$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$

That's an open cover as $C(x, 1/n)^c$ are open.

Take a finite collection $\bigcup_{j=1}^m C(x, 1/n_j)^c = C(x, 1/n_m)^c$

$x \in \bar{K} \Rightarrow C(x, 1/n_m) \cap K \neq \emptyset$.



Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n


$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$

That's an open cover as $C(x, 1/n)^c$ are open.

Take a finite collection $\bigcup_{j=1}^m C(x, 1/n_j)^c = C(x, 1/n_m)^c$

$x \in \bar{K} \Rightarrow C(x, 1/n_m) \cap K \neq \emptyset.$

\Rightarrow so no finite subcover



Suppose K is not closed: $\bar{K} \neq K$

$\Rightarrow \exists x \in \bar{K} \setminus K.$

$x \notin K$ & if $y \neq x \Rightarrow y \in C(x, 1/n)^c$ for some n


$\Rightarrow K \subset \bigcup_{n=1}^{\infty} C(x, 1/n)^c$

That's an open cover as $C(x, 1/n)^c$ are open.

Take a finite collection $\bigcup_{j=1}^m C(x, 1/n_j)^c = C(x, 1/n_m)^c$

$x \in \bar{K} \Rightarrow C(x, 1/n_m) \cap K \neq \emptyset.$

\Rightarrow so no finite subcover $\Rightarrow K$ is not compact.



□

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$.

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K .

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$B(y, 1/n)$

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$B(y, 1/n) \subset B(y, 1/M)$

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$$\begin{aligned} B(y, 1/n) &\subset B(y, 1/M) \\ &\subset B(x, \epsilon/2) \end{aligned}$$

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$$\begin{aligned} B(y, 1/n) &\subset B(y, 1/M) \\ &\subset B(y, \epsilon/2) \subset B(x, \epsilon) \end{aligned}$$

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$$\begin{aligned} B(y, 1/n) &\subset B(y, 1/M) \\ &\subset B(y, \epsilon/2) \subset B(x, \epsilon) \subset U_\lambda, \end{aligned}$$

Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .


Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$$\begin{aligned} B(y, 1/n) &\subset B(y, 1/M) \\ &\subset B(y, \epsilon/2) \subset B(x, \epsilon) \subset U_\lambda, \end{aligned}$$



Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.


U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.

Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$$\begin{aligned} B(y, 1/n) &\subset B(y, 1/M) \\ &\subset B(y, \epsilon/2) \subset B(x, \epsilon) \subset U_\lambda, \end{aligned}$$

($B(y, \epsilon/2) \subset B(x, \epsilon)$ follows by triangle inequality.)



Lemma (Lebesgue covering lemma)

Let (X, d) be a metric space and $K \subset X$. Suppose every sequence in K has a subsequence convergent in K . Given an open cover $\{U_\lambda\}_{\lambda \in I}$ of K , there exists a $\delta > 0$ such that for every $x \in K$, there is a $\lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Proof: If conclusion not true, then \exists open cover $\{U_\lambda\}_{\lambda \in I}$ of K such that:

For every $n \in \mathbb{N}$, $\exists x_n \in K$ s.t. $B(x_n, 1/n)$ is not a subset of any U_λ .

Take $x \in K \Rightarrow \exists \lambda \in I$ s.t. $x \in U_\lambda$.

U_λ is open $\Rightarrow \exists \epsilon > 0$ s.t. $B(x, \epsilon) \subset U_\lambda$.


Take M s.t. $1/M < \epsilon/2$.

If $y \in B(x, \epsilon/2)$ and $n \geq M$, then

$$\begin{aligned} B(y, 1/n) &\subset B(y, 1/M) \\ &\subset B(y, \epsilon/2) \subset B(x, \epsilon) \subset U_\lambda, \end{aligned}$$

($B(y, \epsilon/2) \subset B(x, \epsilon)$ follows by triangle inequality.)

$\Rightarrow y \neq x_n \Rightarrow \{x_n\}_{n=1}^\infty$ has no subsequence converging to x .



Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.
If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

If $n \leq x < n + 1/2$, then $B(x, 1/2) \subset U_{n-1}$.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

If $n \leq x < n + 1/2$, then $B(x, 1/2) \subset U_{n-1}$.

If $n + 1/2 \leq x < n + 1$, then $B(x, 1/2) \subset U_n$.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

If $n \leq x < n + 1/2$, then $B(x, 1/2) \subset U_{n-1}$.

If $n + 1/2 \leq x < n + 1$, then $B(x, 1/2) \subset U_n$.

$\Rightarrow \delta = 1/2$ works.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

If $n \leq x < n + 1/2$, then $B(x, 1/2) \subset U_{n-1}$.

If $n + 1/2 \leq x < n + 1$, then $B(x, 1/2) \subset U_n$.

$\Rightarrow \delta = 1/2$ works.

For the cover given by $U'_n := (\frac{n}{2}, \frac{n+2}{2})$ the largest δ is $1/4$.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

If $n \leq x < n + 1/2$, then $B(x, 1/2) \subset U_{n-1}$.

If $n + 1/2 \leq x < n + 1$, then $B(x, 1/2) \subset U_n$.

$\Rightarrow \delta = 1/2$ works.

For the cover given by $U'_n := (\frac{n}{2}, \frac{n+2}{2})$ the largest δ is $1/4$.

Example: $\mathbb{N} \subset \mathbb{R}$ is not sequentially compact.

Remark: If K such that every sequence has a convergent subsequence in K is called *sequentially compact*.

The δ in the lemma depends on the cover, but not on the x .

Example: $K := [-10, 10]$ and for $n \in \mathbb{Z}$, $U_n := (n, n + 2)$ give an open cover.

If $x \in K$, then $\exists n \in \mathbb{Z}$, such that $n \leq x < n + 1$.

If $n \leq x < n + 1/2$, then $B(x, 1/2) \subset U_{n-1}$.

If $n + 1/2 \leq x < n + 1$, then $B(x, 1/2) \subset U_n$.

$\Rightarrow \delta = 1/2$ works.

For the cover given by $U'_n := (\frac{n}{2}, \frac{n+2}{2})$ the largest δ is $1/4$.

Example: $\mathbb{N} \subset \mathbb{R}$ is not sequentially compact.

Exercise: Find the cover with no $\delta > 0$.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Suppose K is compact and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Suppose K is compact and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

By claim $\exists x \in K$ s.t. $\forall \delta > 0$, $B(x, \delta)$ contains x_n for infinitely many $n \in \mathbb{N}$.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Suppose K is compact and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

By claim $\exists x \in K$ s.t. $\forall \delta > 0$, $B(x, \delta)$ contains x_n for infinitely many $n \in \mathbb{N}$.

$B(x, 1)$ contains some $x_k \Rightarrow$ let $n_1 := k$.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Suppose K is compact and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

By claim $\exists x \in K$ s.t. $\forall \delta > 0$, $B(x, \delta)$ contains x_n for infinitely many $n \in \mathbb{N}$.

$B(x, 1)$ contains some $x_k \Rightarrow$ let $n_1 := k$.

Suppose n_{j-1} is defined.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Suppose K is compact and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

By claim $\exists x \in K$ s.t. $\forall \delta > 0$, $B(x, \delta)$ contains x_n for infinitely many $n \in \mathbb{N}$.

$B(x, 1)$ contains some $x_k \Rightarrow$ let $n_1 := k$.

Suppose n_{j-1} is defined.

$\exists k > n_{j-1}$ such that $x_k \in B(x, 1/j) \Rightarrow$ let $n_j := k$.

Theorem

Let (X, d) be a metric space. Then $K \subset X$ is compact if and only if every sequence in K has a subsequence converging to a point in K .

Proof: Claim: Suppose $K \subset X$ and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K . Suppose $\forall x \in K$, $\exists \alpha_x > 0$ s.t. $x_n \in B(x, \alpha_x)$ for only finitely many $n \in \mathbb{N}$. Then K is not compact.

Proof of the claim: Notice $K \subset \bigcup_{x \in K} B(x, \alpha_x)$.

Any finite subcollection contains x_n for only finitely many n .

\Rightarrow there is no subcover $\Rightarrow K$ is not compact.

Suppose K is compact and $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

By claim $\exists x \in K$ s.t. $\forall \delta > 0$, $B(x, \delta)$ contains x_n for infinitely many $n \in \mathbb{N}$.

$B(x, 1)$ contains some $x_k \Rightarrow$ let $n_1 := k$.

Suppose n_{j-1} is defined.

$\exists k > n_{j-1}$ such that $x_k \in B(x, 1/j) \Rightarrow$ let $n_j := k$.

$d(x, x_{n_j}) < 1/j \Rightarrow \{x_{n_j}\}_{j=1}^{\infty}$ converges to x .

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .

By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Suppose λ_{n-1} is defined.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done),

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

$\exists \lambda_n \in I$ s.t. $B(x_n, \delta) \subset U_{\lambda_n}$.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.


$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.

Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

$\exists \lambda_n \in I$ s.t. $B(x_n, \delta) \subset U_{\lambda_n}$.



Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.


Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

$\exists \lambda_n \in I$ s.t. $B(x_n, \delta) \subset U_{\lambda_n}$.

Either we find a subcover,



Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
 By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.


Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

$\exists \lambda_n \in I$ s.t. $B(x_n, \delta) \subset U_{\lambda_n}$.

Either we find a subcover,
 or we obtain a sequence $\{x_n\}_{n=1}^\infty$
 in K such that $d(x_n, x_k) \geq \delta$
 whenever $k \neq n$.



Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.


Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

$\exists \lambda_n \in I$ s.t. $B(x_n, \delta) \subset U_{\lambda_n}$.

Either we find a subcover,
or we obtain a sequence $\{x_n\}_{n=1}^\infty$
in K such that $d(x_n, x_k) \geq \delta$
whenever $k \neq n$.



No subsequence of $\{x_n\}_{n=1}^\infty$ can be Cauchy.

Now suppose K is sequentially compact and $\{U_\lambda\}_{\lambda \in I}$ is an open cover of K .
By Lebesgue covering lemma, $\exists \delta > 0$ s.t. $\forall x \in K, \exists \lambda \in I$ with $B(x, \delta) \subset U_\lambda$.

Pick $x_1 \in K$ and find $\lambda_1 \in I$ s.t. $B(x_1, \delta) \subset U_{\lambda_1}$.

If $K \subset U_{\lambda_1}$, we are done.

Otherwise, $\exists x_2 \in K \setminus U_{\lambda_1}$. Note $d(x_2, x_1) \geq \delta$.

$\exists \lambda_2 \in I$ such that $B(x_2, \delta) \subset U_{\lambda_2}$. If $K \subset U_{\lambda_1} \cup U_{\lambda_2}$, then done.


Suppose λ_{n-1} is defined.

Either $K \subset U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}}$ (done), or $\exists x_n \in K \setminus (U_{\lambda_1} \cup \dots \cup U_{\lambda_{n-1}})$.

Note $d(x_n, x_j) \geq \delta$ for all $j = 1, 2, \dots, n-1$.

$\exists \lambda_n \in I$ s.t. $B(x_n, \delta) \subset U_{\lambda_n}$.

Either we find a subcover,
or we obtain a sequence $\{x_n\}_{n=1}^\infty$
in K such that $d(x_n, x_k) \geq \delta$
whenever $k \neq n$.



No subsequence of $\{x_n\}_{n=1}^\infty$ can be Cauchy. $\Rightarrow \Leftarrow$

□

Example: Bolzano–Weierstrass theorem for sequences of real numbers:
Every bounded sequence in \mathbb{R} has a convergent subsequence.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

As K is closed, E is closed in $K \iff E$ is closed in X .

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

As K is closed, E is closed in $K \iff E$ is closed in X .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in E .

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

As K is closed, E is closed in $K \iff E$ is closed in X .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in E .

$\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

As K is closed, E is closed in $K \iff E$ is closed in X .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in E .

$\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

$\Rightarrow \exists$ a convergent $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $p \in K$.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

As K is closed, E is closed in $K \iff E$ is closed in X .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in E .

$\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

$\Rightarrow \exists$ a convergent $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $p \in K$.

E is closed $\Rightarrow p \in E$.

Example: Bolzano–Weierstrass theorem for sequences of real numbers:

Every bounded sequence in \mathbb{R} has a convergent subsequence.

\Rightarrow Every sequence in $[a, b] \subset \mathbb{R}$ has a convergent subsequence.

\Rightarrow The limit is also in $[a, b]$.

\Rightarrow A closed bounded interval $[a, b] \subset \mathbb{R}$ is (sequentially) compact.

Proposition

Let (X, d) be a metric space and let $K \subset X$ be compact. If $E \subset K$ is a closed set, then E is compact.

As K is closed, E is closed in $K \iff E$ is closed in X .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in E .

$\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

$\Rightarrow \exists$ a convergent $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $p \in K$.

E is closed $\Rightarrow p \in E$.

$\Rightarrow E$ is (sequentially) compact. □

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general.

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general. The theorem does not hold even for subspaces of \mathbb{R}^n , just in \mathbb{R}^n itself.

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general. The theorem does not hold even for subspaces of \mathbb{R}^n , just in \mathbb{R}^n itself.

Example: In $X = (0, \infty)$ the subset $(0, 1]$ is closed and bounded and **not** compact.

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general. The theorem does not hold even for subspaces of \mathbb{R}^n , just in \mathbb{R}^n itself.

Example: In $X = (0, \infty)$ the subset $(0, 1]$ is closed and bounded and **not** compact.

In general, compact \Rightarrow closed and bounded, **but not vice versa!**

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general. The theorem does not hold even for subspaces of \mathbb{R}^n , just in \mathbb{R}^n itself.

Example: In $X = (0, \infty)$ the subset $(0, 1]$ is closed and bounded and **not** compact.

In general, compact \Rightarrow closed and bounded, **but not vice versa!**

Just for emphasis: **Closed and bounded does not imply compact in general!**

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general. The theorem does not hold even for subspaces of \mathbb{R}^n , just in \mathbb{R}^n itself.

Example: In $X = (0, \infty)$ the subset $(0, 1]$ is closed and bounded and **not** compact.

In general, compact \Rightarrow closed and bounded, **but not vice versa!**

Just for emphasis: **Closed and bounded does not imply compact in general!**

An example complete metric space where closed and bounded is not compact is the space of continuous functions $C([0, 1], \mathbb{R})$.

Theorem (Heine–Borel)

A closed bounded subset $K \subset \mathbb{R}^n$ is compact.

Warning: Heine–Borel only holds for \mathbb{R}^n and not for metric spaces in general. The theorem does not hold even for subspaces of \mathbb{R}^n , just in \mathbb{R}^n itself.

Example: In $X = (0, \infty)$ the subset $(0, 1]$ is closed and bounded and **not** compact.

In general, compact \Rightarrow closed and bounded, **but not vice versa!**

Just for emphasis: **Closed and bounded does not imply compact in general!**

An example complete metric space where closed and bounded is not compact is the space of continuous functions $C([0, 1], \mathbb{R})$.

Exercise: The closed unit ball $C(0, 1)$ in $C([0, 1], \mathbb{R})$ is not compact.

Proof: Suppose $n = 1$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact.

K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact.

K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

$\{y_{k_j}\}_{j=1}^{\infty}$ is also a bounded, so \exists convergent subsequence $\{y_{k_{j_i}}\}_{i=1}^{\infty}$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

$\{y_{k_j}\}_{j=1}^{\infty}$ is also a bounded, so \exists convergent subsequence $\{y_{k_{j_i}}\}_{i=1}^{\infty}$.

$\{x_{k_{j_i}}\}_{i=1}^{\infty}$ is also convergent.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

$\{y_{k_j}\}_{j=1}^{\infty}$ is also a bounded, so \exists convergent subsequence $\{y_{k_{j_i}}\}_{i=1}^{\infty}$.

$\{x_{k_{j_i}}\}_{i=1}^{\infty}$ is also convergent.

Let $x := \lim_{i \rightarrow \infty} x_{k_{j_i}}$ and $y := \lim_{i \rightarrow \infty} y_{k_{j_i}}$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

$\{y_{k_j}\}_{j=1}^{\infty}$ is also a bounded, so \exists convergent subsequence $\{y_{k_{j_i}}\}_{i=1}^{\infty}$.

$\{x_{k_{j_i}}\}_{i=1}^{\infty}$ is also convergent.

Let $x := \lim_{i \rightarrow \infty} x_{k_{j_i}}$ and $y := \lim_{i \rightarrow \infty} y_{k_{j_i}}$.

So $\{(x_{k_{j_i}}, y_{k_{j_i}})\}_{i=1}^{\infty}$ converges to (x, y) .

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

$\{y_{k_j}\}_{j=1}^{\infty}$ is also a bounded, so \exists convergent subsequence $\{y_{k_{j_i}}\}_{i=1}^{\infty}$.

$\{x_{k_{j_i}}\}_{i=1}^{\infty}$ is also convergent.

Let $x := \lim_{i \rightarrow \infty} x_{k_{j_i}}$ and $y := \lim_{i \rightarrow \infty} y_{k_{j_i}}$.

So $\{(x_{k_{j_i}}, y_{k_{j_i}})\}_{i=1}^{\infty}$ converges to (x, y) .

$a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all $k \Rightarrow (x, y) \in B$.

Proof: Suppose $n = 1$.

$K \subset [a, b]$. $[a, b]$ is compact. K is a closed subset of a compact set.

$\Rightarrow K$ is compact.

Now consider $n = 2$. Arbitrary n is an exercise.

If $K \subset \mathbb{R}^2$ is bounded, $\exists B = [a, b] \times [c, d] \subset \mathbb{R}^2$ such that $K \subset B$.

If B is compact, then as K is closed it is also compact.

Let $\{(x_k, y_k)\}_{k=1}^{\infty}$ be a sequence in B .

That is, $a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all k .

By BW, \exists convergent subsequence $\{x_{k_j}\}_{j=1}^{\infty}$.

$\{y_{k_j}\}_{j=1}^{\infty}$ is also a bounded, so \exists convergent subsequence $\{y_{k_{j_i}}\}_{i=1}^{\infty}$.

$\{x_{k_{j_i}}\}_{i=1}^{\infty}$ is also convergent.

Let $x := \lim_{i \rightarrow \infty} x_{k_{j_i}}$ and $y := \lim_{i \rightarrow \infty} y_{k_{j_i}}$.

So $\{(x_{k_{j_i}}, y_{k_{j_i}})\}_{i=1}^{\infty}$ converges to (x, y) .

$a \leq x_k \leq b$ and $c \leq y_k \leq d$ for all $k \Rightarrow (x, y) \in B$.

B is compact $\Rightarrow K$ is compact.



Example: Let (X, d) be a metric space with the discrete metric ($d(x, y) = 1$ if $x \neq y$).

Example: Let (X, d) be a metric space with the discrete metric
($d(x, y) = 1$ if $x \neq y$).

Suppose X is an infinite set. Then

Example: Let (X, d) be a metric space with the discrete metric $(d(x, y) = 1 \text{ if } x \neq y)$.

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.

Example: Let (X, d) be a metric space with the discrete metric ($d(x, y) = 1$ if $x \neq y$).

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.
- (ii) Any $K \subset X$ is closed and bounded.

Example: Let (X, d) be a metric space with the discrete metric ($d(x, y) = 1$ if $x \neq y$).

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.
- (ii) Any $K \subset X$ is closed and bounded.
- (iii) $K \subset X$ is compact $\Leftrightarrow K$ is a finite set.

Example: Let (X, d) be a metric space with the discrete metric ($d(x, y) = 1$ if $x \neq y$).

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.
- (ii) Any $K \subset X$ is closed and bounded.
- (iii) $K \subset X$ is compact $\Leftrightarrow K$ is a finite set.
- (iv) The conclusion of the Lebesgue covering lemma is always satisfied, e.g. with $\delta = 1/2$, even for noncompact $K \subset X$.

Example: Let (X, d) be a metric space with the discrete metric ($d(x, y) = 1$ if $x \neq y$).

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.
- (ii) Any $K \subset X$ is closed and bounded.
- (iii) $K \subset X$ is compact $\Leftrightarrow K$ is a finite set.
- (iv) The conclusion of the Lebesgue covering lemma is always satisfied, e.g. with $\delta = 1/2$, even for noncompact $K \subset X$.

Proof: Exercises.

Example: Let (X, d) be a metric space with the discrete metric $(d(x, y) = 1 \text{ if } x \neq y)$.

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.
- (ii) Any $K \subset X$ is closed and bounded.
- (iii) $K \subset X$ is compact $\Leftrightarrow K$ is a finite set.
- (iv) The conclusion of the Lebesgue covering lemma is always satisfied, e.g. with $\delta = 1/2$, even for noncompact $K \subset X$.

Proof: Exercises.

Remark: Compactness only depends on topology (on the set of open sets).

Example: Let (X, d) be a metric space with the discrete metric $(d(x, y) = 1 \text{ if } x \neq y)$.

Suppose X is an infinite set. Then

- (i) (X, d) is a complete metric space.
- (ii) Any $K \subset X$ is closed and bounded.
- (iii) $K \subset X$ is compact $\Leftrightarrow K$ is a finite set.
- (iv) The conclusion of the Lebesgue covering lemma is always satisfied, e.g. with $\delta = 1/2$, even for noncompact $K \subset X$.

Proof: Exercises.

Remark: Compactness only depends on topology (on the set of open sets).

Completeness depends on which sequences are Cauchy, so it depends on the actual metric.

Exercise: Finite sets are compact.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: A compact set is a complete metric space.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: A compact set is a complete metric space.

Exercise: Suppose (X, d) is complete and $E_1 \supset E_2 \supset E_3 \supset \cdots$ are compact and nonempty. Prove $\bigcap_{j=1}^{\infty} E_j \neq \emptyset$.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: A compact set is a complete metric space.

Exercise: Suppose (X, d) is complete and $E_1 \supset E_2 \supset E_3 \supset \cdots$ are compact and nonempty. Prove $\bigcap_{j=1}^{\infty} E_j \neq \emptyset$.

Exercise: Let (X, d) be a complete metric space. Show that $K \subset X$ is compact if and only if K is closed and such that for every $\epsilon > 0$ there exists a finite set of points x_1, x_2, \dots, x_n with $K \subset \bigcup_{j=1}^n B(x_j, \epsilon)$.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: A compact set is a complete metric space.

Exercise: Suppose (X, d) is complete and $E_1 \supset E_2 \supset E_3 \supset \cdots$ are compact and nonempty. Prove $\bigcap_{j=1}^{\infty} E_j \neq \emptyset$.

Exercise: Let (X, d) be a complete metric space. Show that $K \subset X$ is compact if and only if K is closed and such that for every $\epsilon > 0$ there exists a finite set of points x_1, x_2, \dots, x_n with $K \subset \bigcup_{j=1}^n B(x_j, \epsilon)$. Note: Such a set K is said to be *totally bounded*, so in a complete metric space a set is compact if and only if it is closed and totally bounded.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: A compact set is a complete metric space.

Exercise: Suppose (X, d) is complete and $E_1 \supset E_2 \supset E_3 \supset \cdots$ are compact and nonempty. Prove $\bigcap_{j=1}^{\infty} E_j \neq \emptyset$.

Exercise: Let (X, d) be a complete metric space. Show that $K \subset X$ is compact if and only if K is closed and such that for every $\epsilon > 0$ there exists a finite set of points x_1, x_2, \dots, x_n with $K \subset \bigcup_{j=1}^n B(x_j, \epsilon)$. Note: Such a set K is said to be *totally bounded*, so in a complete metric space a set is compact if and only if it is closed and totally bounded.

Exercise: Prove the general Bolzano–Weierstrass theorem: Any bounded sequence $\{x_k\}_{k=1}^{\infty}$ in \mathbb{R}^n has a convergent subsequence.

Exercise: Finite sets are compact.

Exercise: The union of finitely many compact sets is compact.

Exercise: A compact set is a complete metric space.

Exercise: Suppose (X, d) is complete and $E_1 \supset E_2 \supset E_3 \supset \cdots$ are compact and nonempty. Prove $\bigcap_{j=1}^{\infty} E_j \neq \emptyset$.

Exercise: Let (X, d) be a complete metric space. Show that $K \subset X$ is compact if and only if K is closed and such that for every $\epsilon > 0$ there exists a finite set of points x_1, x_2, \dots, x_n with $K \subset \bigcup_{j=1}^n B(x_j, \epsilon)$. Note: Such a set K is said to be *totally bounded*, so in a complete metric space a set is compact if and only if it is closed and totally bounded.

Exercise: Prove the general Bolzano–Weierstrass theorem: Any bounded sequence $\{x_k\}_{k=1}^{\infty}$ in \mathbb{R}^n has a convergent subsequence.

Exercise: Let (X, d) be a metric space and $K \subset X$. Prove that K is compact as a subset of (X, d) if and only if K is compact as a subset of itself with the subspace metric.