

BA: 7.5

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X$

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Take M such that $\forall n \geq M$, $d_X(x_n, c) < \delta$

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Take M such that $\forall n \geq M$, $d_X(x_n, c) < \delta$

$\Rightarrow \forall n \geq M$, $d_Y(f(x_n), f(c)) < \epsilon$

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Take M such that $\forall n \geq M$, $d_X(x_n, c) < \delta$

$\Rightarrow \forall n \geq M$, $d_Y(f(x_n), f(c)) < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Take M such that $\forall n \geq M$, $d_X(x_n, c) < \delta$

$\Rightarrow \forall n \geq M$, $d_Y(f(x_n), f(c)) < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Now suppose f is not continuous at c .

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Take M such that $\forall n \geq M$, $d_X(x_n, c) < \delta$

$\Rightarrow \forall n \geq M$, $d_Y(f(x_n), f(c)) < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Now suppose f is not continuous at c .

$\exists \epsilon > 0$, s.t. for every $n \in \mathbb{N} \exists x_n \in X$, $d_X(x_n, c) < 1/n$ and $d_Y(f(x_n), f(c)) \geq \epsilon$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and $c \in X$.

$f: X \rightarrow Y$ is *continuous at c* if

$\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $x \in X$ and $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$.

If $f: X \rightarrow Y$ is continuous at all $c \in X$, then f is a *continuous function*.

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces. Then $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every $\{x_n\}_{n=1}^{\infty}$ in X converging to c , $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: Suppose f is continuous at c and $\{x_n\}_{n=1}^{\infty}$ converges to c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $d_X(x, c) < \delta$ implies $d_Y(f(x), f(c)) < \epsilon$.

Take M such that $\forall n \geq M$, $d_X(x_n, c) < \delta$

$\Rightarrow \forall n \geq M$, $d_Y(f(x_n), f(c)) < \epsilon \Rightarrow \{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Now suppose f is not continuous at c .

$\exists \epsilon > 0$, s.t. for every $n \in \mathbb{N} \exists x_n \in X$, $d_X(x_n, c) < 1/n$ and $d_Y(f(x_n), f(c)) \geq \epsilon$.

$\Rightarrow \{x_n\}_{n=1}^{\infty}$ converges to c , but $\{f(x_n)\}_{n=1}^{\infty}$ does not converge to $f(c)$.



Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10} x + a_{01} y + a_{20} x^2 + a_{11} xy + a_{02} y^2 + \cdots + a_{0d} y^d,$$

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.
 $\Rightarrow \lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$.

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.
 $\Rightarrow \lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$.

Then

$$\lim_{n \rightarrow \infty} f(x_n, y_n)$$

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.
 $\Rightarrow \lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$.

Then

$$\lim_{n \rightarrow \infty} f(x_n, y_n) = \lim_{n \rightarrow \infty} \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x_n^j y_n^k$$

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.
 $\Rightarrow \lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$.

Then

$$\lim_{n \rightarrow \infty} f(x_n, y_n) = \lim_{n \rightarrow \infty} \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x_n^j y_n^k = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k$$

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.

$$\Rightarrow \lim_{n \rightarrow \infty} x_n = x \text{ and } \lim_{n \rightarrow \infty} y_n = y.$$

Then

$$\lim_{n \rightarrow \infty} f(x_n, y_n) = \lim_{n \rightarrow \infty} \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x_n^j y_n^k = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = f(x, y).$$

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.
 $\Rightarrow \lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$.

Then

$$\lim_{n \rightarrow \infty} f(x_n, y_n) = \lim_{n \rightarrow \infty} \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x_n^j y_n^k = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = f(x, y).$$

$\Rightarrow f$ is continuous at (x, y)

Example: Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a polynomial:

$$f(x, y) = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = a_{00} + a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \cdots + a_{0d}y^d,$$

Claim: f is continuous.

Let $\{(x_n, y_n)\}_{n=1}^{\infty}$ be a sequence in \mathbb{R}^2 converging to $(x, y) \in \mathbb{R}^2$.
 $\Rightarrow \lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$.

Then

$$\lim_{n \rightarrow \infty} f(x_n, y_n) = \lim_{n \rightarrow \infty} \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x_n^j y_n^k = \sum_{j=0}^d \sum_{k=0}^{d-j} a_{jk} x^j y^k = f(x, y).$$

$\Rightarrow f$ is continuous at $(x, y) \quad \Rightarrow f$ is continuous.


Careful about taking limits separately in \mathbb{R}^n .

Careful about taking limits separately in \mathbb{R}^n .

Consider $f(x, y) := \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$ and $f(0, 0) := 0$.


Careful about taking limits separately in \mathbb{R}^n .

Consider $f(x, y) := \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$ and $f(0, 0) := 0$.



Careful about taking limits separately in \mathbb{R}^n .


Consider $f(x, y) := \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$ and $f(0, 0) := 0$.



Exercise: Prove that f is not continuous at $(0, 0)$,

Careful about taking limits separately in \mathbb{R}^n .


Consider $f(x, y) := \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$ and $f(0, 0) := 0$.



Exercise: Prove that f is not continuous at $(0, 0)$,
but that for every fixed y , the function $g(x) := f(x, y)$ is continuous,

Careful about taking limits separately in \mathbb{R}^n .

Consider $f(x, y) := \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$ and $f(0, 0) := 0$.



Exercise: Prove that f is not continuous at $(0, 0)$,
but that for every fixed y , the function $g(x) := f(x, y)$ is continuous,
and for every fixed x , $h(y) := f(x, y)$ is continuous.

Let X be a metric space and $f: X \rightarrow \mathbb{C}$ a complex-valued function.

Let X be a metric space and $f: X \rightarrow \mathbb{C}$ a complex-valued function.

Write $f(p) = g(p) + i h(p)$, where $g: X \rightarrow \mathbb{R}$ and $h: X \rightarrow \mathbb{R}$.

Let X be a metric space and $f: X \rightarrow \mathbb{C}$ a complex-valued function.

Write $f(p) = g(p) + i h(p)$, where $g: X \rightarrow \mathbb{R}$ and $h: X \rightarrow \mathbb{R}$.

Claim: f is continuous at $c \in X \iff g$ and h are continuous at c .

Let X be a metric space and $f: X \rightarrow \mathbb{C}$ a complex-valued function.

Write $f(p) = g(p) + i h(p)$, where $g: X \rightarrow \mathbb{R}$ and $h: X \rightarrow \mathbb{R}$.

Claim: f is continuous at $c \in X \iff g$ and h are continuous at c .

Claim follows as $\{f(p_n) = g(p_n) + i h(p_n)\}_{n=1}^{\infty}$ converges to $f(p) = g(p) + i h(p)$

Let X be a metric space and $f: X \rightarrow \mathbb{C}$ a complex-valued function.

Write $f(p) = g(p) + i h(p)$, where $g: X \rightarrow \mathbb{R}$ and $h: X \rightarrow \mathbb{R}$.

Claim: f is continuous at $c \in X \iff g$ and h are continuous at c .

Claim follows as $\{f(p_n) = g(p_n) + i h(p_n)\}_{n=1}^{\infty}$ converges to $f(p) = g(p) + i h(p)$

$\iff \{g(p_n)\}_{n=1}^{\infty}$ converges to $g(p)$

Let X be a metric space and $f: X \rightarrow \mathbb{C}$ a complex-valued function.

Write $f(p) = g(p) + i h(p)$, where $g: X \rightarrow \mathbb{R}$ and $h: X \rightarrow \mathbb{R}$.

Claim: f is continuous at $c \in X \iff g$ and h are continuous at c .

Claim follows as $\{f(p_n) = g(p_n) + i h(p_n)\}_{n=1}^{\infty}$ converges to $f(p) = g(p) + i h(p)$

$\iff \{g(p_n)\}_{n=1}^{\infty}$ converges to $g(p)$ and $\{h(p_n)\}_{n=1}^{\infty}$ converges to $h(p)$.

Continuous maps do not map closed sets to closed sets.

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

K is compact $\Rightarrow \exists$ subsequence $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $x \in K$.

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

K is compact $\Rightarrow \exists$ subsequence $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $x \in K$.

By continuity $\lim_{j \rightarrow \infty} f(x_{n_j})$

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

K is compact $\Rightarrow \exists$ subsequence $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $x \in K$.

By continuity $\lim_{j \rightarrow \infty} f(x_{n_j}) = f(x)$

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

K is compact $\Rightarrow \exists$ subsequence $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $x \in K$.

By continuity $\lim_{j \rightarrow \infty} f(x_{n_j}) = f(x) \in f(K)$.

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

K is compact $\Rightarrow \exists$ subsequence $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $x \in K$.

By continuity $\lim_{j \rightarrow \infty} f(x_{n_j}) = f(x) \in f(K)$.

\Rightarrow every sequence in $f(K)$ has a subsequence convergent to a point in $f(K)$

Continuous maps do not map closed sets to closed sets.

E.g., if $f: (0, 1) \rightarrow \mathbb{R}$, $f(x) := x$, then $f((0, 1)) = (0, 1)$,
but $(0, 1)$ is closed in $(0, 1)$, but not closed in \mathbb{R} .

Lemma

*Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$ a continuous function.
If $K \subset X$ is compact, then $f(K)$ is compact.*

Proof: A sequence in $f(K)$ can be written as $\{f(x_n)\}_{n=1}^{\infty}$, where $\{x_n\}_{n=1}^{\infty}$ is a sequence in K .

K is compact $\Rightarrow \exists$ subsequence $\{x_{n_j}\}_{j=1}^{\infty}$ converging to some $x \in K$.

By continuity $\lim_{j \rightarrow \infty} f(x_{n_j}) = f(x) \in f(K)$.

\Rightarrow every sequence in $f(K)$ has a subsequence convergent to a point in $f(K)$

$\Rightarrow f(K)$ is (sequentially) compact. □

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

Proof: X is compact and f continuous

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

Proof: X is compact and f continuous $\Rightarrow f(X) \subset \mathbb{R}$ is compact.

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

Proof: X is compact and f continuous $\Rightarrow f(X) \subset \mathbb{R}$ is compact.

Thus $f(X)$ is closed and bounded.

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

Proof: X is compact and f continuous $\Rightarrow f(X) \subset \mathbb{R}$ is compact.

Thus $f(X)$ is closed and bounded.

$\Rightarrow \sup f(X) \in f(X)$ and $\inf f(X) \in f(X)$

(both achieved by sequences in $f(X)$, which is closed).

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

Proof: X is compact and f continuous $\Rightarrow f(X) \subset \mathbb{R}$ is compact.

Thus $f(X)$ is closed and bounded.

$\Rightarrow \sup f(X) \in f(X)$ and $\inf f(X) \in f(X)$

(both achieved by sequences in $f(X)$, which is closed).

$\Rightarrow \exists x \in X$ s.t. $f(x) = \sup f(X)$

$f: X \rightarrow \mathbb{R}$ achieves an *absolute minimum* at $c \in X$ if

$$f(x) \geq f(c) \quad \text{for all } x \in X.$$

f achieves an *absolute maximum* at $c \in X$ if

$$f(x) \leq f(c) \quad \text{for all } x \in X.$$

Theorem

Let (X, d) be a nonempty compact metric space and let $f: X \rightarrow \mathbb{R}$ be continuous.

$\Rightarrow f$ is bounded and achieves an absolute minimum and an absolute maximum on X .

Proof: X is compact and f continuous $\Rightarrow f(X) \subset \mathbb{R}$ is compact.

Thus $f(X)$ is closed and bounded.

$\Rightarrow \sup f(X) \in f(X)$ and $\inf f(X) \in f(X)$

(both achieved by sequences in $f(X)$, which is closed).

$\Rightarrow \exists x \in X$ s.t. $f(x) = \sup f(X)$ and $\exists y \in X$ s.t. $f(y) = \inf f(X)$.



Lemma


Let (X, d_X) and (Y, d_Y) be metric spaces.

Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Lemma


Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .


Proof: Suppose f is continuous at c , and



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .


Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.
 $\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.




Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,



Lemma


Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.



Lemma


Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,
then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta)$



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .


Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon))$



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .


Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .


Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,
and $B_X(c, \delta)$ is an open neighborhood of c .



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.


$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,
and $B_X(c, \delta)$ is an open neighborhood of c .

Now let $\epsilon > 0$ be given.



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,


then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,

and $B_X(c, \delta)$ is an open neighborhood of c .

Now let $\epsilon > 0$ be given.

If $f^{-1}(B_Y(f(c), \epsilon))$ contains an open neighborhood W of c , then



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.


I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,

and $B_X(c, \delta)$ is an open neighborhood of c .

Now let $\epsilon > 0$ be given.

If $f^{-1}(B_Y(f(c), \epsilon))$ contains an open neighborhood W of c , then

$\exists \delta > 0$, s.t. $B_X(c, \delta) \subset W \subset f^{-1}(B_Y(f(c), \epsilon))$.



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,


and $B_X(c, \delta)$ is an open neighborhood of c .

Now let $\epsilon > 0$ be given.

If $f^{-1}(B_Y(f(c), \epsilon))$ contains an open neighborhood W of c , then

$\exists \delta > 0$, s.t. $B_X(c, \delta) \subset W \subset f^{-1}(B_Y(f(c), \epsilon))$.

That means if $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$,



Lemma

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is continuous at $c \in X \iff$ for every open neighborhood U of $f(c)$ in Y , $f^{-1}(U)$ contains an open neighborhood of c in X .

Proof: Suppose f is continuous at c , and $U \subset Y$ is an open neighborhood of $f(c)$.

$\Rightarrow B_Y(f(c), \epsilon) \subset U$ for some $\epsilon > 0$.

$\exists \delta > 0$, s.t. if $d_X(x, c) < \delta$,

then $d_Y(f(x), f(c)) < \epsilon$.

I.e., $B_X(c, \delta) \subset f^{-1}(B_Y(f(c), \epsilon)) \subset f^{-1}(U)$,

and $B_X(c, \delta)$ is an open neighborhood of c .


Now let $\epsilon > 0$ be given.

If $f^{-1}(B_Y(f(c), \epsilon))$ contains an open neighborhood W of c , then

$\exists \delta > 0$, s.t. $B_X(c, \delta) \subset W \subset f^{-1}(B_Y(f(c), \epsilon))$.

That means if $d_X(x, c) < \delta$, then $d_Y(f(x), f(c)) < \epsilon$,

$\Rightarrow f$ is continuous at c .



□

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces.

$f: X \rightarrow Y$ is continuous \Leftrightarrow for every open $U \subset Y$, $f^{-1}(U)$ is open in X .

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces.

$f: X \rightarrow Y$ is continuous \Leftrightarrow for every open $U \subset Y$, $f^{-1}(U)$ is open in X .

Proof: Exercise.

Example: If $f: X \rightarrow Y$ is continuous and E is closed,

Example: If $f: X \rightarrow Y$ is continuous and E is closed, then $f^{-1}(E) = X \setminus f^{-1}(E^c)$ is also closed.

Example: If $f: X \rightarrow Y$ is continuous and E is closed, then $f^{-1}(E) = X \setminus f^{-1}(E^c)$ is also closed.

Suppose $f: X \rightarrow \mathbb{R}$ is continuous.

Example: If $f: X \rightarrow Y$ is continuous and E is closed, then $f^{-1}(E) = X \setminus f^{-1}(E^c)$ is also closed.

Suppose $f: X \rightarrow \mathbb{R}$ is continuous.

\Rightarrow the zero set of f , $f^{-1}(0) = \{x \in X : f(x) = 0\}$, is closed.

Example: If $f: X \rightarrow Y$ is continuous and E is closed, then $f^{-1}(E) = X \setminus f^{-1}(E^c)$ is also closed.

Suppose $f: X \rightarrow \mathbb{R}$ is continuous.

\Rightarrow the *zero set* of f , $f^{-1}(0) = \{x \in X : f(x) = 0\}$, is closed.

For example, the zero set of a polynomial is closed.

Example: If $f: X \rightarrow Y$ is continuous and E is closed, then $f^{-1}(E) = X \setminus f^{-1}(E^c)$ is also closed.

Suppose $f: X \rightarrow \mathbb{R}$ is continuous.

\Rightarrow the zero set of f , $f^{-1}(0) = \{x \in X : f(x) = 0\}$, is closed.

For example, the zero set of a polynomial is closed.

Similarly,

$f^{-1}([0, \infty)) = \{x \in X : f(x) \geq 0\}$ is closed.

Example: If $f: X \rightarrow Y$ is continuous and E is closed, then $f^{-1}(E) = X \setminus f^{-1}(E^c)$ is also closed.

Suppose $f: X \rightarrow \mathbb{R}$ is continuous.

\Rightarrow the zero set of f , $f^{-1}(0) = \{x \in X : f(x) = 0\}$, is closed.

For example, the zero set of a polynomial is closed.

Similarly,

$f^{-1}([0, \infty)) = \{x \in X : f(x) \geq 0\}$ is closed.

$f^{-1}((0, \infty)) = \{x \in X : f(x) > 0\}$ is open.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

By Lebesgue covering lemma,

$\exists \delta > 0$ s.t. $\forall x \in X, \exists c \in X$ for which $B(x, \delta) \subset B(c, \delta_c)$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

By Lebesgue covering lemma,

$\exists \delta > 0$ s.t. $\forall x \in X, \exists c \in X$ for which $B(x, \delta) \subset B(c, \delta_c)$.

Suppose $p, q \in X$ where $d_X(p, q) < \delta$. Find $c \in X$ s.t. $B(p, \delta) \subset B(c, \delta_c)$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

By Lebesgue covering lemma,

$\exists \delta > 0$ s.t. $\forall x \in X, \exists c \in X$ for which $B(x, \delta) \subset B(c, \delta_c)$.

Suppose $p, q \in X$ where $d_X(p, q) < \delta$. Find $c \in X$ s.t. $B(p, \delta) \subset B(c, \delta_c)$. $\Rightarrow q \in B(c, \delta_c)$.

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

By Lebesgue covering lemma,

$\exists \delta > 0$ s.t. $\forall x \in X, \exists c \in X$ for which $B(x, \delta) \subset B(c, \delta_c)$.

Suppose $p, q \in X$ where $d_X(p, q) < \delta$. Find $c \in X$ s.t. $B(p, \delta) \subset B(c, \delta_c)$. $\Rightarrow q \in B(c, \delta_c)$.

By the triangle inequality and the definition of δ_c ,

$d_Y(f(p), f(q))$

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

By Lebesgue covering lemma,

$\exists \delta > 0$ s.t. $\forall x \in X, \exists c \in X$ for which $B(x, \delta) \subset B(c, \delta_c)$.

Suppose $p, q \in X$ where $d_X(p, q) < \delta$. Find $c \in X$ s.t. $B(p, \delta) \subset B(c, \delta_c)$. $\Rightarrow q \in B(c, \delta_c)$.

By the triangle inequality and the definition of δ_c ,

$$d_Y(f(p), f(q)) \leq d_Y(f(p), f(c)) + d_Y(f(c), f(q))$$

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces. $f: X \rightarrow Y$ is *uniformly continuous* if $\forall \epsilon > 0 \exists \delta > 0$ s.t. whenever $p, q \in X$ and $d_X(p, q) < \delta$, then $d_Y(f(p), f(q)) < \epsilon$.

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces. Suppose $f: X \rightarrow Y$ is continuous and X is compact. Then f is uniformly continuous.

Proof: Let $\epsilon > 0$ be given.

$\forall c \in X$, pick $\delta_c > 0$ s.t. $d_Y(f(x), f(c)) < \epsilon/2$ whenever $x \in B(c, \delta_c)$.

$X \subset \bigcup_{c \in X} B(c, \delta_c)$ and X is compact.

By Lebesgue covering lemma,

$\exists \delta > 0$ s.t. $\forall x \in X, \exists c \in X$ for which $B(x, \delta) \subset B(c, \delta_c)$.

Suppose $p, q \in X$ where $d_X(p, q) < \delta$. Find $c \in X$ s.t. $B(p, \delta) \subset B(c, \delta_c)$. $\Rightarrow q \in B(c, \delta_c)$.

By the triangle inequality and the definition of δ_c ,

$$d_Y(f(p), f(q)) \leq d_Y(f(p), f(c)) + d_Y(f(c), f(q)) < \epsilon/2 + \epsilon/2 = \epsilon.$$



Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous.

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Suppose $|z - y| < \delta$.

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Suppose $|z - y| < \delta$. Then

$$|g(z) - g(y)|$$

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Suppose $|z - y| < \delta$. Then

$$|g(z) - g(y)| = \left| \int_a^b f(x, z) dx - \int_a^b f(x, y) dx \right|$$

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Suppose $|z - y| < \delta$. Then

$$\begin{aligned} |g(z) - g(y)| &= \left| \int_a^b f(x, z) dx - \int_a^b f(x, y) dx \right| \\ &= \left| \int_a^b (f(x, z) - f(x, y)) dx \right| \end{aligned}$$

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Suppose $|z - y| < \delta$. Then

$$\begin{aligned} |g(z) - g(y)| &= \left| \int_a^b f(x, z) dx - \int_a^b f(x, y) dx \right| \\ &= \left| \int_a^b (f(x, z) - f(x, y)) dx \right| \leq (b-a) \frac{\epsilon}{b-a} \end{aligned}$$

Proposition

If $f: [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous, then $g: [c, d] \rightarrow \mathbb{R}$ defined by

$$g(y) := \int_a^b f(x, y) dx \quad \text{is continuous.}$$

Proof: Suppose $y \in [c, d]$.

Let $\epsilon > 0$ be given.

As $[a, b] \times [c, d]$ is compact, f is uniformly continuous. So

$\exists \delta > 0$ s.t. if $z \in [c, d]$ and $|z - y| < \delta$, then $|f(x, z) - f(x, y)| < \frac{\epsilon}{b-a} \quad \forall x \in [a, b]$.

Suppose $|z - y| < \delta$. Then

$$\begin{aligned} |g(z) - g(y)| &= \left| \int_a^b f(x, z) dx - \int_a^b f(x, y) dx \right| \\ &= \left| \int_a^b (f(x, z) - f(x, y)) dx \right| \leq (b-a) \frac{\epsilon}{b-a} = \epsilon. \end{aligned}$$

□

Example: Suppose (X, d_X) and (Y, d_Y) are metric spaces.

Example: Suppose (X, d_X) and (Y, d_Y) are metric spaces.

$f: X \rightarrow Y$ is *Lipschitz* or *K-Lipschitz* if $\exists K \in \mathbb{R}$ s.t.

$$d_Y(f(p), f(q)) \leq K d_X(p, q) \quad \text{for all } p, q \in X.$$

Example: Suppose (X, d_X) and (Y, d_Y) are metric spaces.

$f: X \rightarrow Y$ is *Lipschitz* or *K-Lipschitz* if $\exists K \in \mathbb{R}$ s.t.

$$d_Y(f(p), f(q)) \leq K d_X(p, q) \quad \text{for all } p, q \in X.$$

A Lipschitz f is uniformly continuous:

Example: Suppose (X, d_X) and (Y, d_Y) are metric spaces.

$f: X \rightarrow Y$ is *Lipschitz* or *K-Lipschitz* if $\exists K \in \mathbb{R}$ s.t.

$$d_Y(f(p), f(q)) \leq K d_X(p, q) \quad \text{for all } p, q \in X.$$

A Lipschitz f is uniformly continuous: Take $\delta = \epsilon/K$.

Example: Suppose (X, d_X) and (Y, d_Y) are metric spaces.

$f: X \rightarrow Y$ is *Lipschitz* or *K-Lipschitz* if $\exists K \in \mathbb{R}$ s.t.

$$d_Y(f(p), f(q)) \leq K d_X(p, q) \quad \text{for all } p, q \in X.$$

A Lipschitz f is uniformly continuous: Take $\delta = \epsilon/K$.

f can be uniformly continuous but not Lipschitz:

Example: Suppose (X, d_X) and (Y, d_Y) are metric spaces.

$f: X \rightarrow Y$ is *Lipschitz* or *K-Lipschitz* if $\exists K \in \mathbb{R}$ s.t.

$$d_Y(f(p), f(q)) \leq K d_X(p, q) \quad \text{for all } p, q \in X.$$

A Lipschitz f is uniformly continuous: Take $\delta = \epsilon/K$.

f can be uniformly continuous but not Lipschitz: \sqrt{x} on $[0, 1]$

Definition

Let (X, d) be a metric space and $S \subset X$.

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X, p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function.

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X, p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Suppose $\exists L \in Y$ and $\forall \epsilon > 0, \exists \delta > 0$ s.t. if $x \in S \setminus \{p\}$ and $d_X(x, p) < \delta$, then

$$d_Y(f(x), L) < \epsilon.$$

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X, p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Suppose $\exists L \in Y$ and $\forall \epsilon > 0, \exists \delta > 0$ s.t. if $x \in S \setminus \{p\}$ and $d_X(x, p) < \delta$, then

$$d_Y(f(x), L) < \epsilon.$$

Then $f(x)$ *converges* to L as x goes to p ,

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X, p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Suppose $\exists L \in Y$ and $\forall \epsilon > 0, \exists \delta > 0$ s.t. if $x \in S \setminus \{p\}$ and $d_X(x, p) < \delta$, then

$$d_Y(f(x), L) < \epsilon.$$

Then $f(x)$ *converges* to L as x goes to p , and L is a *limit* of $f(x)$ as x goes to p .

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X, p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Suppose $\exists L \in Y$ and $\forall \epsilon > 0, \exists \delta > 0$ s.t. if $x \in S \setminus \{p\}$ and $d_X(x, p) < \delta$, then

$$d_Y(f(x), L) < \epsilon.$$

Then $f(x)$ *converges* to L as x goes to p , and L is a *limit* of $f(x)$ as x goes to p .

If L is unique, write

$$\lim_{x \rightarrow p} f(x) := L.$$

Definition

Let (X, d) be a metric space and $S \subset X$.

$p \in X$ is a *cluster point* of S if $\forall \epsilon > 0, B(p, \epsilon) \cap S \setminus \{p\} \neq \emptyset$.

Exercise: $p \in X$ is a cluster point of $S \iff p \in \overline{S \setminus \{p\}}$.

Definition

Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X, p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Suppose $\exists L \in Y$ and $\forall \epsilon > 0, \exists \delta > 0$ s.t. if $x \in S \setminus \{p\}$ and $d_X(x, p) < \delta$, then

$$d_Y(f(x), L) < \epsilon.$$

Then $f(x)$ *converges* to L as x goes to p , and L is a *limit* of $f(x)$ as x goes to p .

If L is unique, write

$$\lim_{x \rightarrow p} f(x) := L.$$

If $f(x)$ does not converge as x goes to p , then f *diverges* at p .

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p .

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p . Then the limit of $f(x)$ as x goes to p is unique.

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p . Then the limit of $f(x)$ as x goes to p is unique.

Proof: Exercise.

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p . Then the limit of $f(x)$ as x goes to p is unique.

Proof: Exercise.

Lemma

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function.

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p . Then the limit of $f(x)$ as x goes to p is unique.

Proof: Exercise.

Lemma

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Then $f(x)$ converges to $L \in Y$ as x goes to p \Leftrightarrow for every $\{x_n\}_{n=1}^{\infty}$ in $S \setminus \{p\}$ s.t. $\lim_{n \rightarrow \infty} x_n = p$, $\{f(x_n)\}_{n=1}^{\infty}$ converges to L .

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p . Then the limit of $f(x)$ as x goes to p is unique.

Proof: Exercise.

Lemma

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Then $f(x)$ converges to $L \in Y$ as x goes to p \Leftrightarrow for every $\{x_n\}_{n=1}^{\infty}$ in $S \setminus \{p\}$ s.t. $\lim_{n \rightarrow \infty} x_n = p$, $\{f(x_n)\}_{n=1}^{\infty}$ converges to L .

Proof: Exercise.

Proposition

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function such that $f(x)$ converges as x goes to p . Then the limit of $f(x)$ as x goes to p is unique.

Proof: Exercise.

Lemma

Let (X, d_X) , (Y, d_Y) be metric spaces, $S \subset X$, $p \in X$ a cluster point of S , and $f: S \rightarrow Y$ a function. Then $f(x)$ converges to $L \in Y$ as x goes to p \Leftrightarrow for every $\{x_n\}_{n=1}^{\infty}$ in $S \setminus \{p\}$ s.t. $\lim_{n \rightarrow \infty} x_n = p$, $\{f(x_n)\}_{n=1}^{\infty}$ converges to L .

Proof: Exercise.

Exercise: As on the real numbers, if p is a cluster point of S , then $f: S \rightarrow Y$ is continuous at p if and only if

$$\lim_{x \rightarrow p} f(x) = f(p).$$

Let X and Y be metric spaces.

Let X and Y be metric spaces.

Exercise: If $f: X \rightarrow Y$ is continuous and $A \subset X$, then $f(\overline{A}) \subset \overline{f(A)}$. The subset can be proper.

Let X and Y be metric spaces.

Exercise: If $f: X \rightarrow Y$ is continuous and $A \subset X$, then $f(\overline{A}) \subset \overline{f(A)}$. The subset can be proper.

Exercise: If $f: X \rightarrow Y$ is continuous and X connected, then $f(X)$ is connected.

Let X and Y be metric spaces.

Exercise: If $f: X \rightarrow Y$ is continuous and $A \subset X$, then $f(\bar{A}) \subset \overline{f(A)}$. The subset can be proper.

Exercise: If $f: X \rightarrow Y$ is continuous and X connected, then $f(X)$ is connected.

Exercise: “Intermediate value theorem.” Let (X, d) be connected and $f: X \rightarrow \mathbb{R}$ continuous. If $x_0, x_1 \in X$ and $y \in \mathbb{R}$ such that $f(x_0) < y < f(x_1)$, then $\exists z \in X$ such that $f(z) = y$.

Let X and Y be metric spaces.

Exercise: If $f: X \rightarrow Y$ is continuous and $A \subset X$, then $f(\bar{A}) \subset \overline{f(A)}$. The subset can be proper.

Exercise: If $f: X \rightarrow Y$ is continuous and X connected, then $f(X)$ is connected.

Exercise: “Intermediate value theorem.” Let (X, d) be connected and $f: X \rightarrow \mathbb{R}$ continuous. If $x_0, x_1 \in X$ and $y \in \mathbb{R}$ such that $f(x_0) < y < f(x_1)$, then $\exists z \in X$ such that $f(z) = y$.

Exercise: Suppose $f: X \rightarrow Y$ is one-to-one, onto, and continuous. Suppose X is compact. Then the inverse $f^{-1}: Y \rightarrow X$ is continuous.